arXiv:math/0605379v1 [math.RA] 15 May 2006

AN EXTENDED FREUDENTHAL MAGIC SQUARE IN CHARACTERISTIC 3

ISABEL CUNHA$^\diamond$ AND ALBERTO ELDUQUE*

Abstract. Freudenthal's Magic Square, which in characteristic 0 contains the exceptional Lie algebras other than G_2, is extended over fields of characteristic 3, through the use of symmetric composition superalgebras, to a larger square that contains both Lie algebras and superalgebras. With one exception, the simple Lie superalgebras that appear have no counterpart in characteristic 0.

1. Introduction

In 1966 [Tit66], Tits gave a unified construction of the exceptional simple Lie algebras based on two ingredients: a unital composition algebra (or Hurwitz algebra) and a central simple degree 3 Jordan algebra. This construction is valid over arbitrary fields of characteristic $\neq 2, 3$. The outcome of this construction is given by Freudenthal's Magic Square [Sch66, Fre64] (Table 1).

		dim J			
		6	9	15	27
dim A	1	A_1	A_2	C_3	F_4
	2	A_2	$A_2 \oplus A_2$	A_5	E_6
	4	C_3	A_5	D_6	E_7
	8	F_4	E_6	E_7	E_8

Table 1. Freudenthal's Magic Square

At least in the split cases, this is a construction that depends on two Hurwitz algebras, since the central simple degree Jordan algebras turn out to be the algebras of hermitian 3×3 matrices over a Hurwitz algebra. Even though the construction is not symmetric on the two Hurwitz algebras involved, the result (the Magic Square) is symmetric.

Date: May 15, 2006.

2000 *Mathematics Subject Classification.* Primary 17B25.

Key words and phrases. Freudenthal's Magic Square, simple Lie superalgebras, characteristic 3.

$^\diamond$ Supported by Centro de Matemática da Universidade de Coimbra – FCT.

* Supported by the Spanish Ministerio de Educación y Ciencia and FEDER (MTM 2004-081159-C04-02) and by the Diputación General de Aragón (Grupo de Investigación de Álgebra).

Over the years, several symmetric constructions of Freudenthal's Magic Square based on two Hurwitz algebras have been proposed. Vinberg, in 1966 [Vin05] gave a construction based on two Hurwitz algebras and their derivation algebras. In 1993, Allison and Faulkner [AF93] gave a very general construction based on structurable algebras. In particular, the tensor product of two Hurwitz algebras is structurable and, in this case, the construction is equivalent to the one given by Vinberg. More recently, Barton and Sudbery [BS, BS03] and Landsberg and Manivel [LM02, LM04] provided a different construction based on two Hurwitz algebras and their Lie algebras of triality. (See [Bae02] for a very nice exposition.)

The triality phenomenon was described by E. Cartan [Car38], relating the natural and the two half spin representations of the orthogonal Lie algebra in dimension 8. As shown in [KMRT98], simpler formulas can be obtained if the so called *symmetric composition algebras* are used, instead of the classical Hurwitz algebras. This led the second author to reinterpret the construction by Barton and Sudbery in terms of a construction of a Lie algebra $\mathfrak{g}(S, S')$ out of two symmetric composition algebras S and S' [Eld04]. With a few exceptions in dimension 2, any symmetric composition algebra of dimension 1, 2 or 4 is a *para-Hurwtiz* algebra, while in dimension 8, besides the para-Hurwitz algebras, there appears a new family of symmetric composition algebras, the *Okubo* algebras. The construction in [Eld04] using para-Hurwitz algebras reduces naturally to the construction by Barton and Sudbery (although with slightly simpler formulas). Okubo algebras provide new constructions that highlight different order 3 automorphisms and different distinguished subalgebras of the excepctional simple Lie algebras. Further descriptions of the Lie algebras that appear in Freudenthal's Magic Square in terms of copies of the three-dimensional simple Lie algebra $\mathfrak{sl}_2$ and its two dimensional natural module are derived in [Eldb].

An interesting advantage of the constructions of the Magic Square in terms of two composition algebras and their triality Lie algebras is that this construction is also valid over fields of characteristic 3. Only some attention has to be paid to the second row (or column). Thus [Eld04], for instance, instead of a simple Lie algebra of type E_6, a non simple Lie algebra of dimension 78 is obtained if a two dimensional and an eight dimensional composition algebras are used as ingredients, but it contains a simple codimension 1 ideal of type E_6 (the simple Lie algebra of type E_6 has dimension 77 in characteristic 3).

The characteristic 3 presents also another exceptional feature. Only over fields of this characteristic there are composition superalgebras which are nontrivial (that is, they are not composition algebras). In [EO02] both Hurwitz superalgebras and symmetric composition superalgebras are classified. The main results assert that the only nontrivial superalgebras appear in characteristic 3 and dimensions 3 and 6, in agreement with the classification of the simple alternative superalgebras by Shestakov [She97].

The symmetric composition superalgebras can thus be plugged into the construction $\mathfrak{g}(S, S')$ in [Eld04]. Then an extension in characteristic 3 is obtained of Freudenthal's Magic Square, in which Lie superalgebras appear (see Table 2).

		$\dim S$					
		1	2	4	8	3	6
$\dim S'$	1	A_1	$\tilde{A}_2$	C_3	F_4	$(6,8)$	$(21,14)$
	2		$\tilde{A}_2 \oplus \tilde{A}_2$	$\tilde{A}_5$	$\tilde{E}_6$	$(11,14)$	$(35,20)$
	4			D_6	E_7	$(24,26)$	$(66,32)$
	8				E_8	$(55,50)$	$(133,56)$
	3					$(21,16)$	$(36,40)$
	6						$(78,64)$

Table 2. Freudenthal's Magic Supersquare (characteristic 3)

In Table 2 only the entries above the diagonal have been displayed, as the square is symmetric. The notation $\tilde{X}$ indicates that in characteristic 3 the algebra obtained is not simple, but contains a simple codimension 1 ideal of type X. For instance, in the split case, $\tilde{A}_5$ denotes the projective general Lie algebra $\mathfrak{pgl}_6$. Besides, only the pairs $(\dim \mathfrak{g}_{\bar{0}}, \dim \mathfrak{g}_{\bar{1}})$ are displayed for the nontrivial Lie superalgebras that appear in the "supersquare".

The aim of this paper is the description of all the Lie superalgebras that appear in Table 2. With just a single exception, which corresponds to the pair $(6,8)$, these Lie superalgebras have no counterpart in characteristic 0, and they are either simple or contain a simple codimension 1 ideal (this happens again only in the second row). They will be described as contragredient Lie superalgebras, and to do so, a previous description in terms similar to those used in [Eldb] will be provided too.

In a forthcoming paper, most of these Lie superalgebras will be shown to be isomorphic to the Lie superalgebras constructed in terms of orthogonal and symplectic triple systems in [Eld06b].

The paper is organized as follows. Section 2 will be devoted to review the symmetric composition superalgebras and their Lie superalgebras of triality. Then Section 3 will deal with the construction of a Lie superalgebra in terms of two symmetric composition superalgebras, thus extending the construction in [Eld04]. Section 4 will give the definitions and characterizations of contragredient Lie superalgebras in a way suitable for our purposes. Finally, the last Section 5 will be devoted to the description of all the split Lie superalgebras in Freudenthal's Magic Supersquare as contragredient Lie superalgebras. Their even and odd parts will be computed too.

2. Symmetric composition superalgebras

Recall [EO02] that a *quadratic superform* on a $\mathbb{Z}_2$-graded vector space $U = U_{\bar{0}} \oplus U_{\bar{1}}$ over a field k is a pair $q = (q_{\bar{0}}, b)$ where

(i) $q_{\bar{0}} : U_{\bar{0}} \to k$ is a quadratic form.

(ii) $b : U \times U \to k$ is a supersymmetric even bilinear form. That is, $b|_{U_{\bar 0}\times U_{\bar 0}}$ is symmetric, $b|_{U_{\bar 1}\times U_{\bar 1}}$ is alternating ($b(x_{\bar 1}, x_{\bar 1}) = 0$ for any $x_{\bar 1} \in U_{\bar 1}$) and $b(U_{\bar 0}, U_{\bar 1}) = 0 = b(U_{\bar 1}, U_{\bar 0})$.

(iii) $b|_{U_{\bar 0}\times U_{\bar 0}}$ is the polar of $q_{\bar 0}$. That is, $b(x_{\bar 0}, y_{\bar 0}) = q_{\bar 0}(x_{\bar 0} + y_{\bar 0}) - q_{\bar 0}(x_{\bar 0}) - q_{\bar 0}(y_{\bar 0})$ for any $x_{\bar 0}, y_{\bar 0} \in U_{\bar 0}$.

The quadratic superform $q = (q_{\bar 0}, b)$ is said to be *regular* if $q_{\bar 0}$ is regular (definition as in [KMRT98, p. xix]) and the restriction of b to $U_{\bar 1}$ is nondegenerate.

Then a superalgebra $C = C_{\bar 0} \oplus C_{\bar 1}$ over k, endowed with a regular quadratic superform $q = (q_{\bar 0}, b)$, called the *norm*, is said to be a *composition superalgebra* in case

$$q_{\bar 0}(x_{\bar 0}, y_{\bar 0}) = q_{\bar 0}(x_{\bar 0})q_{\bar 0}(y_{\bar 0}), \tag{2.1a}$$

$$b(x_{\bar 0}y, x_{\bar 0}z) = q_{\bar 0}(x_{\bar 0})b(y, z) = b(yx_{\bar 0}, zx_{\bar 0}), \tag{2.1b}$$

$$b(xy, zt) + (-1)^{|x||y|+|x||z|+|y||z|}b(zy, xt) = (-1)^{|y||z|}b(x, z)b(y, t), \tag{2.1c}$$

for any $x_{\bar 0}, y_{\bar 0} \in C_{\bar 0}$ and homogeneous elements $x, y, z, t \in C$ (where $|x|$ denotes the parity of the homogeneous element x).

The unital composition superalgebras are termed *Hurwitz superalgebras.*

A composition superalgebra $S = S_{\bar 0} \oplus S_{\bar 1}$ with norm $q = (q_{\bar 0}, b)$ is said to be *symmetric* if

$$b(xy, z) = b(x, yz),$$

for any $x, y, z \in S$. That is, the associated supersymmetric bilinear form is associative.

Example 2.2. Let V be a two dimensional vector space over the field k with a nonzero alternating bilinear form $\langle .|. \rangle$. Consider the superspace

$$B(1, 2) = k1 \oplus V,$$

where $B(1, 2)_{\bar 0} = k1$ and $B(1, 2)_{\bar 1} = V$, with supercommutative multiplication given by

$$1x = x1 = x \qquad \text{and} \qquad uv = \langle u|v \rangle 1$$

for any $x \in B(1, 2)$ and $u, v \in V$, and with quadratic superform $q = (q_{\bar 0}, b)$ given by:

$$q_{\bar 0}(1) = 1, \quad b(u, v) = \langle u|v \rangle,$$

for any $u, v \in V$.

If the characteristic of k is 3, then $B(1, 2)$ is a Hurwitz superalgebra [EO02, Proposition 2.7].

Example 2.3. Let V be as in Example 2.2, then $\mathrm{End}_k(V)$ is equipped with the symplectic involution $f \mapsto \bar f$, such that

$$\langle f(u)|v \rangle = \langle u|\bar f(v) \rangle,$$

for any $u, v \in V$. Consider the superspace

$$B(4, 2) = \mathrm{End}_k(V) \oplus V,$$

where $B(4, 2)_{\bar 0} = \mathrm{End}_k(V)$ and $B(4, 2)_{\bar 1} = V$, with multiplication given by:

- the usual multiplication (composition of maps) in $\mathrm{End}_k(V)$,

- $v\cdot f = f(v) = \bar f\cdot v$ for any $f\in \operatorname{End}_k(V)$ and $v\in V$,
- $u\cdot v = \langle .|u\rangle v\ \bigl(w\mapsto \langle w|u\rangle v\bigr)\ \in \operatorname{End}_k(V)$ for any $u,v\in V$,

and with quadratic superform $q=(q_{\bar 0},b)$ such that

$$q_{\bar 0}(f)=\det f,\quad b(u,v)=\langle u|v\rangle,$$

for any $f\in\operatorname{End}_k(V)$ and $u,v\in V$.

Again, if the characteristic is 3, $B(4,2)$ is a Hurwitz superalgebra [EO02, Proposition 2.7].

The vector space $\operatorname{End}_k(V)$ may be identified with $V\otimes V$ (unadorned tensor products are assumed to be tensor products over the ground field k) by means of

$$\begin{aligned} V\otimes V &\longrightarrow \operatorname{End}_k(V)\\ x\otimes y &\mapsto \langle x|.\rangle y : v\mapsto \langle x|v\rangle y. \end{aligned}$$

The symplectic involution on $\operatorname{End}_k(V)$ becomes now $\overline{x\otimes y}=-y\otimes x$ for any $x,y\in V$.

Then $B(4,2)$ can be identified with

$$B(4,2)=(V\otimes V)\oplus V, \tag{2.4}$$

with multiplication given by

$$\begin{cases} (x\otimes y)\cdot(z\otimes t)=\langle x|t\rangle z\otimes y,\\ u\cdot(x\otimes y)=\langle x|u\rangle y=-(y\otimes x)\cdot u\\ u\cdot v=-u\otimes v, \end{cases}$$

for any $x,y,z,t,u,v\in V$, and the supersymmetric bilinear form b is determined by

$$b(x\otimes y,z\otimes t)=\langle x|z\rangle\langle y|t\rangle,\quad b(u,v)=\langle u|v\rangle, \tag{2.5}$$

for any $x,y,z,t,u,v\in V$ (see [Eldb, §2]).

Given any Hurwitz superalgebra C with norm $q=(q_{\bar 0},b)$, the linear map $x\mapsto \bar x=b(x,1)1-x$ is a superinvolution ($\bar{\bar x}=x$ and $\overline{xy}=(-1)^{|x||y|}\bar y\bar x$ for any homogeneous $x,y\in C$). Then C, with the same norm q, but with new multiplication

$$x\bullet y=\bar x\bar y$$

becomes a symmetric composition superalgebra, which is called a *para-Hurwitz superalgebra*, and denoted by $\bar C$.

Also, as for the ungraded case, if φ is an order 3 automorphism of a symmetric composition superalgebra S, with multiplication $\bullet$ and norm q (we will refer to this as the symmetric composition superalgebra $(S,\bullet,q)$), then φ is an isometry of q, and with the new multiplication given by

$$x*y=\varphi(x)\bullet\varphi^2(y), \tag{2.6}$$

$(S,*,q)$ is again a symmetric composition algebra, denoted by S_φ or $(S,\bullet,q)_\varphi$.

Example 2.7. Given any order 3 automorphism of $B(1,2)$ over a field k of characteristic 3 (and hence this is also an automorphism of the associated para-Hurwitz superalgebra $\overline{B(1,2)}$), there is a scalar $\lambda \in k$ and a symplectic basis $\{v, w\}$ of V (that is, $\langle v|w\rangle = 1$) such that $\varphi(v) = v$ and $\varphi(w) = \lambda v + w$. The symmetric composition superalgebra defined on $B(1,2)$ by means of the new multiplication given by

$$x \bullet y = \varphi(\bar{x})\varphi^2(\bar{y})$$

is denoted by $B(1,2)_\lambda$ (see [EO02, Example 2.9]).

For $\lambda = 1$ this is just the para-Hurwitz superalgebra $\overline{B(1,2)}$.

Example 2.8. In terms of the description of $B(4,2)$ as $(V \otimes V) \oplus V$, the multiplication in the associated para-Hurwtiz superalgebra $\overline{B(4,2)}$ is determined by:

$$\begin{cases} (x \otimes y) \bullet (z \otimes t) = \langle y|z\rangle t \otimes x, \\ u \bullet (x \otimes y) = \langle y|u\rangle x = -(y \otimes x) \bullet u \\ u \bullet v = -u \otimes v, \end{cases}$$

for any $x, y, z, t, u, v \in V$. The associated bilinear form is given in (2.5).

Remark 2.9. This is exactly the multiplication in the para-Cayley algebra in [Eldb, eq. (2.2)], if the last copy of V is suppressed there.

The classification of the symmetric composition superalgebras appears in [EO02, Theorem 4.3]. Over fields of characteristic 2, any symmetric composition superalgebra is a symmetric composition algebra suitably graded over $\mathbb{Z}_2$. In characteristic $\neq 2$, the classification is given by:

Theorem 2.10. *Let k be a field of characteristic $\neq 2$, and let S be a symmetric composition superalgebras over k. Then either:*

(i) $S_{\bar{1}} = 0$*; that is, S is a symmetric composition algebra, or*

(ii) *the characteristic of k is 3 and either there is a scalar $\lambda \in k$ such that S is isomorphic to $\overline{B(1,2)}_\lambda$ or S is isomorphic to $\overline{B(4,2)}$.*

Therefore, the superalgebras in Examples 2.7 and 2.8 exhaust, up to isomorphism, all the symmetric composition superalebras, which do not appear in the ungraded setting.

In order to superize the construction of the Lie algebras $\mathfrak{g}(S, S')$ in the Introduction, the triality Lie superalgebras are needed. Given a symmetric composition superalgebra $(S, *, q)$, its *triality Lie superalgebra* $\mathfrak{tri}(S, *, q) = \mathfrak{tri}(S, *, q)_{\bar{0}} \oplus \mathfrak{tri}(S, *, q)_{\bar{1}}$ is defined by

$$\begin{aligned} &\mathfrak{tri}(S, *, q)_i \\ &\quad = \bigl\{(d_0, d_1, d_2) \in \mathfrak{osp}(S, q)_i^3 : d_0(x * y) = d_1(x) * y + (-1)^{i|x|} x * d_2(y) \\ &\qquad\qquad \text{for any homogeneous elements } x, y \in S\bigr\}, \end{aligned}$$

where $i = \bar{0}, \bar{1}$. For simplicity, this Lie superalgebra will be referred to simply by $\mathfrak{tri}(S)$. Here $\mathfrak{osp}(S, q)$ denotes the orthosymplectic Lie superalgebra of the superform q. The bracket in $\mathfrak{tri}(S)$ is given componentwise.

Note that $\mathfrak{tri}(S)$ is endowed with a natural automorphism $\theta : (d_0, d_1, d_2) \mapsto (d_2, d_0, d_1)$, satisfying $\theta^3 = 1$.

Theorem 2.11. *Let $(S, *, q)$ be any of the symmetric composition superalgebras $\overline{B(1,2)}$ or $\overline{B(4,2)}$. Then for any homogeneous element $d_0 \in \mathfrak{osp}(S,q)_i$ ($i = \bar{0}, \bar{1}$), there are unique $d_1, d_2 \in \mathfrak{osp}(S,q)_i$ such that $(d_0, d_1, d_2) \in \mathfrak{tri}(S)$. Besides, the map $\Phi : \mathfrak{osp}(S,q) \to \mathfrak{osp}(S,q)$ given by $\Phi(d_0) = d_1$ is an automorphism of $\mathfrak{osp}(S,q)$ with $\Phi^3 = 1$. Hence $\mathfrak{tri}(S) = \{\big(d, \Phi(d), \Phi^2(d)\big) : d \in \mathfrak{osp}(S,q)\}$ is isomorphic to $\mathfrak{osp}(S,q)$. Moreover, for $\overline{B(1,2)}$, this automorphism Φ is the identity.*

Proof. This is [EO02, Theorem 5.6]. The last assertion is proved in [EO02, p. 5464]. ☐

Corollary 2.12. $\mathfrak{tri}(\overline{B(1,2)}) = \{(d,d,d) : d \in \mathfrak{osp}(B(1,2), q)\}$.

Given a nondegenerate even supersymmetric bilinear form b on a superspace $W = W_{\bar{0}} \oplus W_{\bar{1}}$, its orthosymplectic Lie superalgebra $\mathfrak{osp}(W,b)$ is spanned by the operators

$$\sigma_{x,y} : z \mapsto (-1)^{|y||z|} b(x,z)y - (-1)^{|x|(|y|+|z|)} b(y,z)x,$$

for homogeneous $x, y, z \in W$. Besides, for homogeneous $x, y, z, t \in W$,

$$[\sigma_{x,y}, \sigma_{z,t}] = \sigma_{\sigma_{x,y}(z),t} + (-1)^{(|x|+|y|)|z|} \sigma_{z,\sigma_{x,y}(t)}. \tag{2.13}$$

Also, given a vector space U endowed with a nondegenerate alternating bilinear form $\langle .|. \rangle$, its symplectic Lie algebra $\mathfrak{sp}\big(U, \langle .|. \rangle\big)$ is spanned by the operators

$$\gamma_{x,y} : z \mapsto \langle x|z \rangle y + \langle y|z \rangle x, \tag{2.14}$$

for $x, y, z \in U$.

Over any field k of characteristic 3, consider the para-Hurwitz superalgebra $\overline{B(1,2)}$ which, for simplicity, will be denoted by $S_{1,2}$. Thus, $S_{1,2} = k1 \oplus V$, where V is a two dimensional vector space endowed with a nonzero (hence nondegenerate) alternating bilinear form $\langle .|. \rangle$. The norm $q = (q_{\bar{0}}, b)$ satisfies $q_{\bar{0}}(1) = 1$ (so $b(1,1) = 2 = -1$) and $b(u,v) = \langle u|v \rangle$ for any $u, v \in V$. The corresponding orthosymplectic Lie superalgebra $\mathfrak{osp}(S_{1,2}, b) = \sigma_{S_{1,2},S_{1,2}}$ satisfies:

$$\mathfrak{osp}(S_{1,2}, b)_{\bar{0}} = \sigma_{V,V}, \quad \mathfrak{osp}(S_{1,2}, b) = \sigma_{1,V}$$

(note that $\sigma_{1,1} = 0$). But for any $u, v, w \in V$:

$$\sigma_{u,v} : \begin{cases} 1 \mapsto 0, \\ w \mapsto -b(u,w)v - b(v,w)u = -\gamma_{u,v}(w), \end{cases} \tag{2.15}$$

while

$$\sigma_{1,u} : \begin{cases} 1 \mapsto 2u = -u \ (\operatorname{char} k = 3), \\ v \mapsto -b(u,v)1 = -\langle u|v \rangle 1. \end{cases} \tag{2.16}$$

Also, $[\sigma_{1,u}, \sigma_{1,v}] = -\sigma_{u,v}$ and $[\sigma_{u,v}, \sigma_{1,w}] = \sigma_{1,\sigma_{u,v}(w)}$ for any $u, v, w \in V$. Thus, $\sigma_{V,V}$ can be identified with the three dimensional symplectic

Lie algebra $\mathfrak{sp}(V)$, $\sigma_{1,V}$ with V ($\sigma_{1,u} \leftrightarrow u$), and therefore $\mathfrak{osp}(S_{1,2}, b)$ is isomorphic to the Lie superalgebra

$$\mathfrak{b}_{0,1} = \mathfrak{sp}(V) \oplus V, \tag{2.17}$$

with even part $\mathfrak{sp}(V)$, odd part V, and multiplication determined by:

- the Lie algebra $\mathfrak{sp}(V)$ is a subalgebra,
- $[\gamma, v] = \gamma(v)$ for any $\gamma \in \mathfrak{sp}(V)$ and $v \in V$,
- $[u, v] = \gamma_{u,v}$ for any $u, v \in V$.

Now, consider the superalgebra $S_{4,2} = \overline{B(4,2)}$ over a field k of characteristic 3, as described in Example 2.8. Then [EO02, Lemma 5.7] shows that the Lie superalgebra $\mathfrak{osp}(S_{4,2}, b)$ is isomorphic to the Lie superalgebra

$$\mathfrak{d}_{2,1} = \bigl(\mathfrak{sp}(V) \oplus \mathfrak{sp}(V) \oplus \mathfrak{sp}(V)\bigr) \oplus (V \otimes V \otimes V), \tag{2.18}$$

with even part $\mathfrak{sp}(V) \oplus \mathfrak{sp}(V) \oplus \mathfrak{sp}(V)$, odd part $V \otimes V \otimes V$, and multiplication determined by:

- the Lie algebra $\mathfrak{sp}(V) \oplus \mathfrak{sp}(V) \oplus \mathfrak{sp}(V)$ is a subalgebra,
- for any $\gamma_1, \gamma_2, \gamma_3 \in \mathfrak{sp}(V)$ and $u_1, u_2, u_3 \in V$,

$$\begin{aligned}
&[(\gamma_1, \gamma_2, \gamma_3), u_1 \otimes u_2 \otimes u_3] \\
&\qquad = \gamma_1(u_1) \otimes u_2 \otimes u_3 + u_1 \otimes \gamma_2(u_2) \otimes u_3 + u_1 \otimes u_2 \otimes \gamma(u_3),
\end{aligned}$$

- for any $u_1, u_2, u_3, v_1, v_2, v_3 \in V$,

$$\begin{aligned}
&[u_1 \otimes u_2 \otimes u_3, v_1 \otimes v_2 \otimes v_3] \\
&\quad = -\bigl(\langle u_2 | v_2\rangle \langle u_3 | v_3\rangle \gamma_{u_1,v_1}, \langle u_1 | v_1\rangle \langle u_3 | v_3\rangle \gamma_{u_2,v_2}, \langle u_1 | v_1\rangle \langle u_2 | v_2\rangle \gamma_{u_3,v_3}\bigr) \\
&\quad = -\sum_{i=1}^{3} \bigl(\prod_{j \neq i} \langle u_j | v_j\rangle\bigr) \nu_i(\gamma_{u_i,v_i}),
\end{aligned}$$

where $\nu_i : \mathfrak{sp}(V) \to \mathfrak{sp}(V)^3$ denotes the inclusion on the i^{th}-component.

(A word of caution is needed here: the operators $\sigma_{a,b}$ in [EO02] are changed in sign with respect to ours.)

Moreover, the action of $\mathfrak{d}_{2,1}$ on $S_{4,2} = (V \otimes V) \oplus V$ is given by the isomorphism:

$$\rho : \mathfrak{d}_{2,1} \to \mathfrak{osp}(S_{4,2}, b) \tag{2.19}$$

such that

$$\rho\bigl((\gamma_1, \gamma_2, \gamma_3)\bigr)(v_1 \otimes v_2 + v_3) = \bigl(\gamma_1(v_1) \otimes v_2 + v_1 \otimes \gamma_2(v_2)\bigr) + \gamma_3(v_3),$$

$$\begin{aligned}
\rho\bigl(u_1 \otimes u_2 \otimes u_3\bigr)(v_1 \otimes v_2 + v_3) &= -\sigma_{u_1 \otimes u_2, u_3}(v_1 \otimes v_2 + v_3) \\
&= \langle u_3 | v_3\rangle u_1 \otimes u_2 - \langle u_1 | v_1\rangle \langle u_2 | v_2\rangle u_3,
\end{aligned}$$

for any $\gamma_1, \gamma_2, \gamma_3 \in \mathfrak{sp}(V)$ and $u_1, u_2, u_3, v_1, v_2, v_3 \in V$.

Note that $\sigma_{u_3,v_3}(w_3) = -b(u_3, w_3)v_3 - b(v_3, w_3)u_3 = -\gamma_{u_3,v_3}(w_3)$, so (see [Eldb, (2.8)])

$$\begin{aligned}
\sigma_{u_1 \otimes u_2, v_1 \otimes v_2} &= -\rho\bigl(\langle u_2 | v_2\rangle \gamma_{u_1,v_1}, \langle u_1 | v_1\rangle \gamma_{u_2,v_2}, 0)\bigr), \\
\sigma_{u_3,v_3} &= -\rho\bigl((0, 0, \gamma_{u_3,v_3})\bigr), \\
\sigma_{u_1 \otimes u_2, u_3} &= -\rho\bigl(u_1 \otimes u_2 \otimes u_3\bigr),
\end{aligned} \tag{2.20}$$

for any $u_1, u_2, u_3, v_1, v_2, v_3 \in V$.

Consider the natural order 3 automorphism θ of $\mathfrak{d}_{2,1}$ such that

$$\begin{cases} \theta(\gamma_1, \gamma_2, \gamma_3) = (\gamma_3, \gamma_1, \gamma_2), \\ \theta(u_1 \otimes u_2 \otimes u_3) = u_3 \otimes u_1 \otimes u_2, \end{cases} \tag{2.21}$$

for any $\gamma_1, \gamma_2, \gamma_3 \in \mathfrak{sp}(V)$ and $u_1, u_2, u_3 \in V$.

Then (compare to [Eldb, Proposition 2.12]):

Proposition 2.22. *For any homogeneous elements $f \in \mathfrak{d}_{2,1}$ and $x, y \in S_{4,2}$:*

$$\rho(f)(x \bullet y) = \rho(\theta^{-1}(f))(x) \bullet y + (-1)^{|f||x|} x \bullet \rho(\theta^{-2}(f))(y).$$

Proof. It is enough to prove this for generators of $\mathfrak{d}_{2,1}$ and a spanning set of $S_{4,2}$, and hence for $f = u_1 \otimes u_2 \otimes u_3$, $x = v_1 \otimes v_2$ or $x = v_3$, and $y = w_1 \otimes w_2$ or $y = w_3$, where $u_i, v_i, w_i \in V$, $i = 1, 2, 3$. This is straightforward. For instance,

$$\begin{aligned}
\rho(u_1 \otimes u_2 \otimes u_3)\bigl(v_3 \bullet (w_1 \otimes w_2)\bigr) &= \rho(u_1 \otimes u_2 \otimes u_3)\bigl(\langle w_2|v_3\rangle w_1\bigr) \\
&= \langle w_2|v_3\rangle\langle u_3|w_1\rangle u_1 \otimes u_2, \\
\rho\bigl(\theta^{-1}(u_1 \otimes u_2 \otimes u_3)\bigr)(v_3) \bullet (w_1 \otimes w_2) &= \rho(u_2 \otimes u_3 \otimes u_1)(v_3) \bullet (w_1 \otimes w_2) \\
&= \langle u_1|v_3\rangle (u_2 \otimes u_3) \bullet (w_1 \otimes w_2) \\
&= \langle u_1|v_3\rangle\langle u_3|w_1\rangle w_2 \otimes u_2 \\
-v_3 \bullet \rho\bigl(\theta^{-2}(u_1 \otimes u_2 \otimes u_3)\bigr)(w_1 \otimes w_2) &= -v_3 \bullet \bigl(\rho(u_3 \otimes u_1 \otimes u_2)(w_1 \otimes w_2) \\
&= v_3 \bullet \bigl((\langle u_3|w_1\rangle\langle u_1|w_2\rangle u_2 \\
&= -\langle u_3|w_1\rangle\langle u_1|w_2\rangle v_3 \otimes u_2,
\end{aligned}$$

and now, since $\langle u|v\rangle w + \langle v|w\rangle u + \langle w|u\rangle v = 0$ for any $u, v, w \in V$ (as any trilinear alternating form on a two dimensional vector space is trivially zero), one gets

$$\begin{aligned}
\rho(u_1 \otimes u_2 \otimes u_3)\bigl(v_3 \bullet (w_1 \otimes w_2)\bigr) = \rho\bigl(\theta^{-1}(u_1 \otimes u_2 \otimes u_3)\bigr)(v_3) \bullet (w_1 \otimes w_2) \\
- v_3 \bullet \rho\bigl(\theta^{-2}(u_1 \otimes u_2 \otimes u_3)\bigr)(w_1 \otimes w_2),
\end{aligned}$$

as required. ☐

Write $\rho_i = \rho \circ \theta^{-i}$, then Theorem 2.11 and Proposition 2.22 give, as in [Eldb, Corollary 2.13]:

Corollary 2.23. $\mathfrak{tri}(S_{4,2}) = \bigl\{\bigl(\rho_0(f), \rho_1(f), \rho_2(f)\bigr) : f \in \mathfrak{d}_{2,1}\bigr\}$.

This Corollary allows us to identify $\mathfrak{tri}(S_{4,2})$ to $\mathfrak{d}_{2,1}$ by means of $f \leftrightarrow \bigl(\rho_0(f), \rho_1(f), \rho_2(f)\bigr)$.

Remark 2.24. The formulas for the bracket in $\mathfrak{d}_{2,1}$ and for the representations ρ_i of $\mathfrak{d}_{2,1}$ ($i = 0, 1, 2$) are exactly the ones that appear in [Eldb, §2] for $\mathfrak{d}_4$, but with the last copy of V suppressed.

Denote by V_i ($i = 1, 2, 3$) the module V for $\mathfrak{sp}(V)^3 = \mathfrak{sp}(V) \oplus \mathfrak{sp}(V) \oplus \mathfrak{sp}(V)$ on which only the i^{th} component of $\mathfrak{sp}(V)^3$ acts: $(s_1, s_2, s_3).v_i = s_i(v_i)$, for any $s_j \in \mathfrak{sp}(V)$, $j = 1, 2, 3$, and $v_i \in V_i$. Also denote by $\iota_i(S_{4,2})$ the $\mathfrak{d}_{2,1}$-module associated to the representation ρ_i. Then, as modules for $\mathfrak{sp}(V)^3$ (compare to [Eldb, (2.14)]):

$$\begin{cases} \iota_0(S_{4,2}) = (V_1 \otimes V_2) \oplus V_3, \\ \iota_1(S_{4,2}) = (V_2 \otimes V_3) \oplus V_1, \\ \iota_2(S_{4,2}) = (V_3 \otimes V_1) \oplus V_2 \big(\simeq (V_1 \otimes V_3) \oplus V_2\big). \end{cases} \tag{2.25}$$

The multiplication $\bullet$ on $S_{4,2}$ (see Example 2.8) becomes the bilinear $\mathfrak{tri}(S_{4,2})$-invariant map

$$\iota_0(S_{4,2}) \times \iota_1(S_{4,2}) \to \iota_2(S_{4,2})$$

given by,

$$\begin{split} \big((V_1 \otimes V_2) \oplus V_3\big) \times \big(V_2 \otimes V_3) \oplus V_1\big) &\to (V_3 \otimes V_1) \oplus V_2 \\ \big(u_1 \otimes u_2 + u_3), (v_2 \otimes v_3 + v_1\big) &\mapsto \big(\langle u_2|v_2\rangle v_3 \otimes u_1 - u_3 \otimes v_1\big) \\ &\qquad - \big(\langle u_1|v_1\rangle u_2 + \langle u_3|v_3\rangle v_2\big) \end{split} \tag{2.26}$$

for any $u_i, v_i \in V_i$, $i = 1, 2, 3$; and cyclically. Note that it consists of contractions for repeated indices.

This is exactly the multiplication in [Eldb, 3.1] if V_4 is ignored there.

3. The extended Freudenthal's Magic Square

Let $(S, \bullet, q)$ and $(S', *, q')$ be two symmetric composition superalgebras and define $\mathfrak{g} = \mathfrak{g}(S, S')$ to be the $\mathbb{Z}_2 \times \mathbb{Z}_2$-graded anticommutative superalgebra such that

$$\begin{split} \mathfrak{g}_{(\bar 0,\bar 0)} &= \mathfrak{tri}(S, \bullet, q) \oplus \mathfrak{tri}(S', *, q'), \\ \mathfrak{g}_{(\bar 1,\bar 0)} &= \mathfrak{g}_{(\bar 0,\bar 1)} = \mathfrak{g}_{(\bar 1,\bar 1)} = S \otimes S'. \end{split}$$

For any $x \in S$ and $x' \in S'$, denote by $\iota_i(x \otimes x')$ the element $x \otimes x'$ in $\mathfrak{g}_{(\bar 1,\bar 0)}$ (respectively $\mathfrak{g}_{(\bar 0,\bar 1)}$, $\mathfrak{g}_{(\bar 1,\bar 1)}$) if $i = 0$ (respectively, $i = 1, 2$). Thus

$$\mathfrak{g} = \mathfrak{g}(S, S') = \big(\mathfrak{tri}(S, \bullet, q) \oplus \mathfrak{tri}(S', *, q')\big) \oplus \big(\oplus_{i=0}^2 \iota_i(S \otimes S')\big). \tag{3.1}$$

Then $\mathfrak{g}$ is a superalgebra where, for $i = \bar 0, \bar 1$,

$$\mathfrak{g}_i = (\mathfrak{g}_{(\bar 0,\bar 0)})_i \oplus (\mathfrak{g}_{(\bar 1,\bar 0)})_i \oplus (\mathfrak{g}_{(\bar 0,\bar 1)})_i \oplus (\mathfrak{g}_{(\bar 1,\bar 1)})_i,$$

with

$$\begin{split} (\mathfrak{g}_{(\bar 0,\bar 0)})_i &= \mathfrak{tri}(S)_i \oplus \mathfrak{tri}(S')_i, \\ \iota_j(S \otimes S')_i &= \iota_j(S_{\bar 0} \otimes S'_i) \oplus \iota_j(S_{\bar 1} \otimes S'_{\bar 1 - i}), \end{split}$$

for $j = 0, 1, 2$.

The superanticommutative multiplication on $\mathfrak{g}$ is defined by means of:

- $\mathfrak{g}_{(\bar 0,\bar 0)}$ is a Lie subsuperalgebra of $\mathfrak{g}$,
- $[(d_0, d_1, d_2), \iota_i(x \otimes x')] = \iota_i\big(d_i(x) \otimes x'\big)$,
- $[(d'_0, d'_1, d'_2), \iota_i(x \otimes x')] = (-1)^{|d'_i||x|}\iota_i\big(x \otimes d'_i(x')\big)$,
- $[\iota_i(x \otimes x'), \iota_{i+1}(y \otimes y')] = (-1)^{|x'||y|}\iota_{i+2}\big((x \bullet y) \otimes (x' * y')\big)$ (indices modulo 3),

- $$[\iota_i(x\otimes x'),\iota_i(y\otimes y')] = (-1)^{|x||x'|+|x||y'|+|y||y'|}b'(x',y')\theta^i(t_{x,y}) \\ +(-1)^{|y||x'|}b(x,y)\theta'^i(t'_{x',y'}),$$

for any $i=0,1,2$ and homogeneous $x,y\in S$, $x',y'\in S'$, $(d_0,d_1,d_2)\in \mathfrak{tri}(S)$, and $(d'_0,d'_1,d'_2)\in\mathfrak{tri}(S')$. Here θ denotes the natural automorphism $\theta:(d_0,d_1,d_2)\mapsto(d_2,d_0,d_1)$ in $\mathfrak{tri}(S)$, θ' the analogous automorphism of $\mathfrak{tri}(S')$, and

$$t_{x,y}=\bigl(\sigma_{x,y},\tfrac12 b(x,y)1-r_xl_y,\tfrac12 b(x,y)1-l_xr_y\bigr) \tag{3.2}$$

(with $l_x(y)=x\bullet y$, $r_x(y)=(-1)^{|x||y|}y\bullet x$), while $t'_{x',y'}$ is the analogous triple in $\mathfrak{tri}(S')$.

Just superizing the arguments in [Eld04, Theorem 3.1]; that is, taking into account the parity signs, one gets:

Theorem 3.3. *With this multiplication, $\mathfrak{g}=\mathfrak{g}(S,S')$ is a Lie superalgebra.*

With the same proof as in the ungraded case [Eld06a, Theorem 12.2] the following result is obtained:

Proposition 3.4. *Let S and S' be two symmetric composition superalgebras, and let φ be an automorphism of S of order 3, then the Lie superalgebras $\mathfrak{g}(S,S')$ and $\mathfrak{g}(S_\varphi,S')$ are isomorphic.*

(The superalgebra S_φ was defined in (2.6).)

Therefore, over fields of characteristic 3, there is no need to deal with the symmetric composition superalgebras $\overline{B(1,2)}_\lambda$, but just with $S_{1,2}=\overline{B(1,2)}$ (which is three dimensional) and with $S_{4,2}=\overline{B(4,2)}$ (whose dimension is 6). Freudenthal's Magic Square thus extends over these fields to the larger square

		dim S					
		1	2	4	8	3	6
dim S'	1	A_1	$\tilde{A}_2$	C_3	F_4	$\mathfrak{g}(S_1,S_{1,2})$	$\mathfrak{g}(S_1,S_{4,2})$
	2		$\tilde{A}_2\oplus\tilde{A}_2$	$\tilde{A}_5$	$\tilde{E}_6$	$\mathfrak{g}(S_2,S_{1,2})$	$\mathfrak{g}(S_2,S_{4,2})$
	4			D_6	E_7	$\mathfrak{g}(S_4,S_{1,2})$	$\mathfrak{g}(S_4,S_{4,2})$
	8				E_8	$\mathfrak{g}(S_8,S_{1,2})$	$\mathfrak{g}(S_8,S_{4,2})$
	3					$\mathfrak{g}(S_{1,2},S_{1,2})$	$\mathfrak{g}(S_{1,2},S_{4,2})$
	6						$\mathfrak{g}(S_{4,2},S_{4,2})$

(see Table 2, where the dimensions of the even and odd part of the superalgebras involved, which are easily computed from the definitions, are displayed), where S_n denotes a symmetric composition algebra of dimension n ($n=1,2,4$ or 8).

The purpose for the remaining part of the paper is the description of the Lie superalgebras $\mathfrak{g}(S,S_{1,2})$ and $\mathfrak{g}(S,S_{4,2})$ in the split case, that is, in case the (super)algebra S contains nontrivial idempotents (for instance, over an algebraically closed field). Proposition 3.4, or [Eld06a, Corollary 12.1] shows that it is enough to take then for S_n the unique split para-Hurwitz algebras (dimensions 1, 2, 4 or 8), together with $S_{1,2}$ and $S_{4,2}$.

This description will be given in terms of contragredient Lie superalgebras.

4. Contragredient Lie superalgebras

Let $A = \bigl(a_{ij}\bigr)$ be a square matrix of size n over a ground field k, and let $(\mathfrak{h}, \Pi, \Pi^\vee)$ be a *realization* of A, as defined in [Kac90, §1.1]. That is,

- $\Pi = \{\alpha_1, \ldots, \alpha_n\}$ is a linearly independent set in $\mathfrak{h}^*$ (the dual of the vector space $\mathfrak{h}$),
- $\Pi^\vee = \{h_1, \ldots, h_n\}$ is a linearly independent set in $\mathfrak{h}$,
- $\alpha_j(h_i) = a_{ij}$ for any $i, j = 1, \ldots, n$,
- $\dim \mathfrak{h} = 2n - \operatorname{rank} A$.

As in [Kac77], for any subset $\tau \subseteq \{1, \ldots, n\}$, consider the local Lie superalgebra

$$\hat{\mathfrak{g}}(A, \tau) = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1,$$

with $\mathfrak{g}_0 = \mathfrak{h}$, $\mathfrak{g}_1 = ke_1 \oplus \cdots \oplus ke_n$, $\mathfrak{g}_{-1} = kf_1 \oplus \cdots \oplus kf_n$, and bracket given by:

$$[e_i, f_j] = \delta_{ij} h_i, \quad [h, h'] = 0, \quad [h, e_i] = \alpha_i(h) e_i, \quad [h, f_i] = -\alpha_i(h) f_i, \quad (4.1)$$

for any $i, j = 1, \ldots, n$ and $h, h' \in \mathfrak{h}$, where $\mathfrak{g}_0$ is even and e_i, f_i are even if and only if $i \notin \tau$.

Note that changing f_i by cf_i, $0 \neq c \in k$, the i^{th} row of A is multiplied by c. Hence, if $a_{ii} \neq 0$, then a_{ii} can be taken to be 2, as it is customary.

Then (see [Kac77, 1.2.2]) there exists a minimal $\mathbb{Z}$-graded Lie superalgebra $\mathfrak{g}(A, \tau)$ with local part $\hat{\mathfrak{g}}(A, \tau)$. To define $\mathfrak{g}(A, \tau)$ one first considers $\tilde{\mathfrak{g}}(A, \tau)$, the free Lie superalgebra generated by $\mathfrak{h}$ and $\{e_1, \ldots, e_n, f_1, \ldots, f_n\}$, subject to the relations in (4.1), endowed with the $\mathbb{Z}$-grading induced by the grading in $\hat{\mathfrak{g}}(A, \tau)$ (note that the relations are homogeneous), and then considers the largest homogeneous ideal $\mathfrak{i}(A)$ that intersects $\mathfrak{h}$ trivially. Then

$$\mathfrak{g}(A, \tau) = \tilde{\mathfrak{g}}(A, \tau)/\mathfrak{i}(A).$$

Remark 4.2.

- As in [Kac90, Proposition 1.6], with $\mathfrak{g}'(A, \tau) = [\mathfrak{g}(A, \tau), \mathfrak{g}(A, \tau)]$, the center is given by
  $$\mathfrak{c} = Z\bigl(\mathfrak{g}(A, \tau)\bigr) = Z\bigl(\mathfrak{g}'(A, \tau)\bigr) = \{h \in \mathfrak{h} : \alpha_i(h) = 0 \ \forall i = 1, \ldots, n\}.$$
  Note that $\dim \mathfrak{c} = n - \operatorname{rank} A$.
- Let Q be the free abelian group on generators $\epsilon_1, \ldots, \epsilon_n$. Then $\mathfrak{g}(A, \tau)$ is Q-graded by assigning $\deg e_i = \epsilon_i = -\deg f_i$, $i = 1, \ldots, n$, and $\deg h = 0$ for any $h \in \mathfrak{h}$. Now $\tilde{\mathfrak{g}}(A, \tau)$ is Q-graded too, because the relation in (4.1) are Q-homogeneous. For any $m \in \mathbb{Z}$, the m^{th} homogeneous component of $\tilde{\mathfrak{g}}(A, \tau)$ equals
  $$\bigoplus_{\substack{q = m_1\epsilon_1 + \cdots + m_n\epsilon_n \\ m_1 + \cdots + m_n = m}} \tilde{\mathfrak{g}}(A, \tau)_q,$$

and the ideal $\mathfrak{i}(A)$ is Q-homogeneous too, since $\mathfrak{i}(A)$ is contained in $\oplus_{0\neq q\in Q}\pi_q\bigl(\mathfrak{i}(A)\bigr)$ (π_q denotes the projection onto the q^{th}-component), which is a Q-homogeneous ideal that intersects trivially $\mathfrak{h}$. Hence, by maximality, $\mathfrak{i}(A) = \oplus_{0\neq q\in Q}\pi_q\bigl(\mathfrak{i}(A)\bigr)$.

The following notation will be used:

◇ $\mathfrak{g}(A,\tau)$ will be called the *contragredient Lie superalgebra* with Cartan matrix A.

◇ $\mathfrak{g}'(A,\tau)$ will be called the *derived contragredient Lie superalgebra* with Cartan matrix A.

◇ $\mathfrak{g}(A,\tau)/\mathfrak{c}$ will be called the *centerless contragredient Lie superalgebra* with Cartan matrix A.

◇ $\mathfrak{g}'(A,\tau)/\mathfrak{c}$ will be called the *centerless derived contragredient Lie superalgebra* with Cartan matrix A.

A bit of caution is needed here, in [VK71] it is the Lie algebra $\mathfrak{g}'(A,\tau)$ the one that is called the contragredient Lie algebra.

Note that, as in [Kac90, Proposition 1.7], $\mathfrak{g}(A,\tau)$ has no Q-homogeneous ideal if and only if $\det A\neq 0$ and for any i,j there exist indices $i_1,\dots,i_s$ such that $a_{ii_1}a_{i_1i_2}\cdots a_{i_sj}\neq 0$. If only this last condition holds, then any Q-homogeneous ideal either contains $\mathfrak{g}'(A,\tau)$ or is contained in the center. Also note that if $\det A\neq 0$, then $\mathfrak{c}=0$, $\mathfrak{g}'(A,\tau)=\mathfrak{g}(A,\tau)$ and the four Lie superalgebras considered above coincide.

Lemma 4.3. *Under the conditions above, let* $\mathfrak{j}(A)$ *be the largest* $\mathbb{Z}$*-homogeneous ideal of* $\tilde{\mathfrak{g}}(A,\tau)$ *which intersects trivially* $\mathfrak{g}_{-1}\oplus\mathfrak{g}_1$*. Then* $\mathfrak{j}(A) = \mathfrak{i}(A)\oplus\mathfrak{c}$.

Proof. It is clear that $\mathfrak{i}(A)\cap\mathfrak{g}_{\pm1}=0$, otherwise $\mathfrak{i}(A)$ would contain some $0\neq h\in\mathfrak{h}$ (because of $[e_i,f_i]=h_i$ for any i). Hence $\mathfrak{i}(A)$ is the largest homogeneous ideal intersecting trivially $\mathfrak{g}_{-1}\oplus\mathfrak{g}_0\oplus\mathfrak{g}_1$. Thus $\bigl(\mathfrak{i}(A)\oplus\mathfrak{c}\bigr)\cap\bigl(\mathfrak{g}_{-1}\oplus\mathfrak{g}_1\bigr)=0$, so $\mathfrak{i}(A)\oplus\mathfrak{c}\subseteq\mathfrak{j}(A)$.

Conversely, $\tilde{\mathfrak{g}}(A,\tau)=\oplus_{i\in\mathbb{Z}}\tilde{\mathfrak{g}}_i$, with $\tilde{\mathfrak{g}}_i=\mathfrak{g}_i$ for $i=-1,0,1$, and for any $h\in\mathfrak{j}(A)\cap\tilde{\mathfrak{g}}_0=\mathfrak{j}(A)\cap\mathfrak{h}$, $[h,e_i]=\alpha_i(h)e_i\in\mathfrak{j}(A)\cap\mathfrak{g}_1=0$ for any i. Thus $\alpha_i(h)=0$ for any i and $h\in\mathfrak{c}$. Therefore $\mathfrak{j}(A)\subseteq\mathfrak{c}\oplus\bigl(\oplus_{i\neq-1,0,1}\mathfrak{j}(A)\cap\tilde{\mathfrak{g}}_i\bigr)$. But $\oplus_{i\neq-1,0,1}\mathfrak{j}(A)\cap\tilde{\mathfrak{g}}_i$ is an ideal of $\tilde{\mathfrak{g}}(A,\tau)$, since it is closed under the action of e_i, f_i, $i=1,\dots,n$, and $\mathfrak{h}$ (note that $[f_i,\mathfrak{j}(A)\cap\tilde{\mathfrak{g}}_2]\subseteq\mathfrak{j}(A)\cap\mathfrak{g}_1=0$). Hence $\oplus_{i\neq-1,0,1}\mathfrak{j}(A)\cap\tilde{\mathfrak{g}}_i\subseteq\mathfrak{i}(A)$, and $\mathfrak{j}(A)\subseteq\mathfrak{i}(A)\oplus\mathfrak{c}$. □

Let us consider the natural characterizations of the contragredient Lie superalgebras, that will be used in the next section. The previous notations and assumptions will be kept in the following results.

Theorem 4.4. *Let* $\mathfrak{g}$ *be a* $\mathbb{Z}$*-graded Lie superalgebra generated by* $\mathfrak{g}_0$*, which is contained in* $\mathfrak{g}_{\bar 0}$*, and by elements* $e_i\in\mathfrak{g}_1$ *and* $f_i\in\mathfrak{g}_{-1}$*,* $i=1,\dots,n$*, which are even (respectively odd) if and only if* $i\notin\tau$ *(resp.* $i\in\tau$*). Assume that there are* $\alpha_1,\dots,\alpha_n\in\mathfrak{g}_0^*$ *and* $h_1,\dots,h_n\in\mathfrak{g}_0$ *such that*

$$\bigl(\mathfrak{g}_0,\{\alpha_1,\dots,\alpha_n\},\{h_1,\dots,h_n\}\bigr)$$

is a realization of A *and that relations* (4.1) *hold. If any nonzero homogeneous ideal of* $\mathfrak{g}$ *intersects* $\mathfrak{g}_0$ *nontrivially, then* $\mathfrak{g}$ *is isomorphic to the contragredient Lie superalgebra* $\mathfrak{g}(A,\tau)$.

Proof. This is clear from the definitions, which give an epimorphism $\phi : \mathfrak{g}(A,\tau) \to \mathfrak{g}$, such that $\ker\phi \cap \mathfrak{g}(A,\tau)_0 = 0$, so $\ker\phi = 0$. □

Theorem 4.5. *Let* $\mathfrak{g}$ *be a* $\mathbb{Z}$*-graded Lie superalgebra such that:*

- (i) $\mathfrak{g}_0 = \bar{\mathfrak{h}}$ *is contained in* $\mathfrak{g}_{\bar{0}}$*, it is abelian and its dimension is* n*.*
- (ii) *There are linearly independent elements* $\bar{\alpha}_1,\dots,\bar{\alpha}_n \in \bar{\mathfrak{h}}^*$ *and elements* $\bar{h}_1,\dots,\bar{h}_n \in \bar{\mathfrak{h}}$ *such that* $\bar{\alpha}_j(\bar{h}_i) = a_{ij}$ *for any* i,j*.*
- (iii) *There are elements* $\bar{e}_1,\dots,\bar{e}_n \in \mathfrak{g}_1$ *and* $\bar{f}_1,\dots,\bar{f}_n \in \mathfrak{g}_{-1}$*, where* $\bar{e}_i,\bar{f}_i$ *are even (resp. odd) if* $i \notin \tau$ *(resp.* $i \in \tau$*), and such that the relations* (4.1) *are satisfied (with* e_i *replaced by* $\bar{e}_i$*, ...), and* $\mathfrak{g}$ *is generated by* $\bar{e}_1,\dots,\bar{e}_n,\bar{f}_1,\dots,\bar{f}_n$ *and* $\bar{\mathfrak{h}}$*.*
- (iv) *Any nonzero homogeneous ideal of* $\mathfrak{g}$ *intersects* $\mathfrak{g}_{-1}\oplus\mathfrak{g}_1$ *nontrivially.*

Then $\mathfrak{g}$ *is isomorphic to the centerless contragredient Lie superalgebra* $\mathfrak{g}(A,\tau)/\mathfrak{c}$*.*

Proof. The elements $\alpha_1,\dots,\alpha_n \in \mathfrak{h}^*$ ($\mathfrak{h} = \mathfrak{g}(A,\tau)_0$) define linearly independent elements $\hat{\alpha}_i \in (\mathfrak{h}/\mathfrak{c})^*$, because $\mathfrak{c} = \{h \in \mathfrak{h} : \alpha_i(h) = 0\ \forall i\}$. Hence there is a linear bijection $\varphi^* : \bar{\mathfrak{h}}^* \to (\mathfrak{h}/\mathfrak{c})^*$ such that $\varphi^*(\bar{\alpha}_i) = \hat{\alpha}_i$ for any i. Besides, φ^* induces a linear bijection $\varphi : \mathfrak{h}/\mathfrak{c} \to \bar{\mathfrak{h}}$, such that $\bar{\alpha}_j\bigl(\varphi(h+\mathfrak{c})\bigr) = \hat{\alpha}_j(h+\mathfrak{c}) = \alpha_j(h)$ for any j and $h \in \mathfrak{h}$. By composing with the natural projection, a surjective linear map is obtained $\phi : \mathfrak{h} \to \bar{\mathfrak{h}}$ such that $\bar{\alpha}_j\bigl(\phi(h)\bigr) = \alpha_j(h)$ for any j and $h \in \mathfrak{h}$.

Now, by definition of $\tilde{\mathfrak{g}}(A,\tau)$, ϕ extends to a surjective homomorphism of Lie superalgebras $\tilde{\phi} : \tilde{\mathfrak{g}}(A,\tau) \to \mathfrak{g}$ which takes e_i (respectively f_i) to $\bar{e}_i$ (respectively $\bar{f}_i$), for any i, and $\tilde{\phi}$ is homogeneous. Condition (iv) shows that $\tilde{\phi}\bigl(\mathfrak{j}(A)\bigr) = 0$, so $\tilde{\phi}$ induces a surjective homomorphism $\hat{\phi} : \tilde{\mathfrak{g}}(A,\tau)/\mathfrak{j}(A) \to \mathfrak{g}$. But $\hat{\phi}$ is bijective in degrees 0, 1 and -1 (the elements $\bar{e}_1,\dots,\bar{e}_n$ are linearly independent, as they are eigenvectors of eigenvalues $\bar{\alpha}_1,\dots,\bar{\alpha}_n$ for the action of $\bar{\mathfrak{h}}$). By definition of $\mathfrak{j}(A)$, one concludes that $\ker\hat{\phi} = 0$, so $\mathfrak{g}$ is isomorphic to $\tilde{\mathfrak{g}}(A,\tau)/\mathfrak{j}(A)$, which is isomorphic to $\mathfrak{g}(A,\tau)/\mathfrak{c}$ by Lemma 4.3. □

Theorem 4.6. *Let* $\mathfrak{g}$ *be a* Q*-graded Lie superalgebra generated by nonzero elements* $\bar{e}_i \in \mathfrak{g}_{\epsilon_i}$*,* $\bar{f}_i \in \mathfrak{g}_{-\epsilon_i}$*,* $i = 1,\dots,n$*, where* $\bar{e}_i,\bar{f}_i$ *are even (resp. odd) if* $i \notin \tau$ *(resp.* $i \in \tau$*). Assume that:*

- (i) *The elements* $\bar{h}_i = [\bar{e}_i,\bar{f}_i]$*,* $i = 1,\dots,n$*, are linearly independent.*
- (ii) $[\bar{h}_i,\bar{e}_j] = a_{ij}\bar{e}_j$*,* $[\bar{h}_i,\bar{f}_j] = -a_{ij}\bar{f}_j$*,* $[\bar{e}_i,\bar{f}_j] = \delta_{ij}\bar{h}_i$*, and* $[\bar{h}_i,\bar{h}_j] = 0$*, for any* $i,j = 1,\dots,n$*.*
- (iii) *Any nonzero* Q*-homogeneous ideal of* $\mathfrak{g}$ *intersects nontrivially* $\mathfrak{g}_{-1}\oplus\mathfrak{g}_0\oplus\mathfrak{g}_1$*, where* $\mathfrak{g}_m = \oplus\{\tilde{\mathfrak{g}}_q : q = m_1\epsilon_1+\dots+m_n\epsilon_n,\ m_1+\dots+m_n = m\}$ *for any* $m \in \mathbb{Z}$*.*

Then $\mathfrak{g}$ *is isomorphic to the derived contragredient Lie superalgebra* $\mathfrak{g}'(A,\tau)$*.*

Proof. Since $\mathfrak{g}$ is generated by the elements $\bar{e}_1,\dots,\bar{e}_n,\bar{f}_1,\dots,\bar{f}_n$, it follows that $\mathfrak{g}_0 = k\bar{h}_1 \oplus\dots\oplus k\bar{h}_n$. Take an even vector space $\hat{\mathfrak{g}}_0$ of dimension

$n - \operatorname{rank} A$ and elements $\bar\alpha_1, \dots, \bar\alpha_n \in (\mathfrak{g}_0 \oplus \hat{\mathfrak{g}}_0)^*$ such that

$$\begin{cases} \bar\alpha_j(\bar h_i) = a_{ij} \text{ for any } i, j, \\ \bar\alpha_1, \dots, \bar\alpha_n \text{ are linearly independent.} \end{cases}$$

Then $\bigl(\mathfrak{g}_0 \oplus \hat{\mathfrak{g}}_0, \{\bar\alpha_1, \dots, \bar\alpha_n\}, \{\bar h_1, \dots, \bar h_n\}\bigr)$ is a realization of A, and $\bar{\mathfrak{g}} = \mathfrak{g} \oplus \hat{\mathfrak{g}}_0$ is a Q-graded Lie superalgebra , with $\bar{\mathfrak{g}}_0 = \mathfrak{g}_0 \oplus \hat{\mathfrak{g}}_0$, $\bar{\mathfrak{g}}_\epsilon = \mathfrak{g}_\epsilon$ for any $0 \neq \epsilon \in Q$, and where $[\bar{\mathfrak{g}}_0, \bar{\mathfrak{g}}_0] = 0$, $\mathfrak{g}$ is an ideal of $\bar{\mathfrak{g}}$, and for any $h \in \hat{\mathfrak{g}}_0$ and $x \in \mathfrak{g}_\epsilon$ $(0 \neq \epsilon = m_1\epsilon_1 + \dots + m_n\epsilon_n \in Q)$, $[h, x] = (m_1\bar\alpha_1 + \dots + m_n\bar\alpha_n)(h)x$.

By Theorem 4.4, $\bar{\mathfrak{g}}$ is isomorphic to the contragredient Lie superalgebra $\mathfrak{g}(A, \tau)$, and hence $\mathfrak{g} = [\bar{\mathfrak{g}}, \bar{\mathfrak{g}}]$ is isomorphic to $\mathfrak{g}'(A, \tau)$. □

Theorem 4.7. *Let $\mathfrak{g}$ be a Q-graded Lie superalgebra generated by nonzero elements $\bar e_i \in \mathfrak{g}_{\epsilon_i}$, $\bar f_i \in \mathfrak{g}_{-\epsilon_i}$, $i = 1, \dots, n$, where $\bar e_i, \bar f_i$ are even (resp. odd) if $i \notin \tau$ (resp. $i \in \tau$). Assume that:*

- (i) *If $\bar h_i = [\bar e_i, \bar f_i]$, $i = 1, \dots, n$, then $[\bar h_i, \bar e_j] = a_{ij}\bar e_j$, $[\bar h_i, \bar f_j] = -a_{ij}\bar f_j$, $[\bar h_i, \bar h_j] = 0$, and $[\bar e_i, \bar f_j] = \delta_{ij}\bar h_i$, for any i, j.*
- (ii) *Any nonzero Q-homogeneous ideal intersects nontrivially $\mathfrak{g}_{-1} \oplus \mathfrak{g}_1$, where $\mathfrak{g}_m = \oplus\{\tilde{\mathfrak{g}}_q : q = m_1\epsilon_1 + \dots + m_n\epsilon_n,\ m_1 + \dots + m_n = m\}$ for any $m \in \mathbb{Z}$.*

Then $\mathfrak{g}$ is isomorphic to the centerless derived contragredient Lie superalgebra $\mathfrak{g}'(A, \tau)/\mathfrak{c}$.

Proof. Because of item (i), $\mathfrak{g}_0 = \operatorname{span}\bigl\{\bar h_1, \dots, \bar h_n\bigr\}$ holds, and condition (ii) implies that the dimension of $\mathfrak{g}_0$ equals the rank of A (any linear combination of the rows of A which gives 0 induces a linear combination of the $\bar h_i$'s which is central, and hence spans an ideal with trivial intersection with $\mathfrak{g}_{-1} \oplus \mathfrak{g}_1$).

As in the proof of Theorem 4.6 take a vector space $\hat{\mathfrak{g}}_0$ of dimension $n - \operatorname{rank} A$, and elements $\bar\alpha_1, \dots, \bar\alpha_n \in (\mathfrak{g}_0 \oplus \hat{\mathfrak{g}}_0)^*$ such that

$$\begin{cases} \bar\alpha_j(\bar h_i) = a_{ij} \text{ for any } i, j, \\ \bar\alpha_1, \dots, \bar\alpha_n \text{ are linearly independent.} \end{cases}$$

The Lie superalgebra $\bar{\mathfrak{g}}$ constructed as in the proof of Theorem 4.6 satisfies now the hypotheses of Theorem 4.5. Hence $\bar{\mathfrak{g}}$ is isomorphic to the centerless contragredient Lie superalgebra $\mathfrak{g}(A, \tau)/\mathfrak{c}$, and hence $\mathfrak{g} = [\bar{\mathfrak{g}}, \bar{\mathfrak{g}}]$ is isomorphic to the centerless derived contragredient Lie superalgebra $\mathfrak{g}'(A, \tau)/\mathfrak{c} = [\mathfrak{g}(A, \tau)/\mathfrak{c}, \mathfrak{g}(A, \tau)/\mathfrak{c}]$. □

In order to present the Lie superalgebras $\mathfrak{g}(S, S')$ in the Extended Freudenthal's Magic Square as contragredient Lie superalgebras, it will be useful first to describe them in terms similar to those used in [Eldb].

Let V be, as before, a two dimensional vector space endowed with a nonzero alternating bilinear form $\langle .|. \rangle$. For any $n \in \mathbb{N}$ consider copies $V_1, \dots, V_n$ of V, and for any subset $\sigma \subseteq \{1, \dots, n\}$ take the module for $\mathfrak{sp}(V)^n = \mathfrak{sp}(V_1) \oplus \dots \oplus \mathfrak{sp}(V_n)$ given by:

$$V(\sigma) = \begin{cases} \mathfrak{sp}(V_1) \oplus \dots \oplus \mathfrak{sp}(V_n) & \text{if } \sigma = \emptyset, \\ V_{i_1} \otimes \dots \otimes V_{i_r} & \text{if } \sigma = \{i_1, \dots, i_r\},\ 1 \leq i_1 < \dots < i_r \leq n. \end{cases} \tag{4.8}$$

(V_i is the natural module for $\mathfrak{sp}(V_i)$ annihilated by $\mathfrak{sp}(V_j)$ for $j \neq i$.)

Identify, as in [Eldb, Section 2], any subset $\sigma \subseteq \{1,\dots,n\}$ with the element $(\sigma_1,\dots,\sigma_n) \in \mathbb{Z}_2^n$, such that $\sigma_i = 1$ if and only if $i \in \sigma$. Then for any $\sigma, \tau \in \mathbb{Z}_2^n$, consider the natural $\mathfrak{sp}(V)^n$-invariant map:

$$\varphi_{\sigma,\tau} : V(\sigma) \times V(\tau) \longrightarrow V(\sigma+\tau) \tag{4.9}$$

defined as follows:

- If $\sigma \neq \tau$ and $\sigma \neq \emptyset \neq \tau$, then $\varphi_{\sigma,\tau}$ is obtained by contraction, by means of $\langle .|.\rangle$ in the indices $i \in \sigma \cap \tau$ ($\sigma_i = 1 = \tau_i$). Thus, for instance,
$$\varphi_{\{1,2,3\},\{1,3,4\}}(u_1 \otimes u_2 \otimes u_3, v_1 \otimes v_3 \otimes v_4) = \langle u_1|v_1\rangle\langle u_3|v_3\rangle u_2 \otimes v_4$$
for any $u_1, v_1 \in V_1$, $u_2 \in V_2$, $u_3, v_3 \in V_3$ and $v_4 \in V_4$.
- $\varphi_{\emptyset,\emptyset}$ is the Lie bracket in $\mathfrak{sp}(V)^n = \mathfrak{sp}(V_1) \oplus \cdots \oplus \mathfrak{sp}(V_n)$.
- For any $\sigma \neq \emptyset$, $\varphi_{\emptyset,\sigma} = -\varphi_{\sigma,\emptyset}$ is given by the natural action of the Lie algebra $\mathfrak{sp}(V)^n$ on $V(\sigma)$. Thus, for instance,
$$\varphi_{\emptyset,\{1,3\}}\bigl((s_1,\dots,s_n), u_1 \otimes u_3\bigr) = s_1(u_1) \otimes u_3 + u_1 \otimes s_3(u_3),$$
for any $s_i \in \mathfrak{sp}(V)$, $i = 1,\dots,n$, and $u_1 \in V_1$, $u_3 \in V_3$.
- Finally, for any $\sigma = \{i_1,\dots,i_r\} \neq \emptyset$, $\varphi_{\sigma,\sigma}$ is given by:
$$\varphi_{\sigma,\sigma}(u_{i_1} \otimes \cdots \otimes u_{i_r}, v_{i_1} \otimes \cdots \otimes v_{i_r}) = \sum_{j=1}^{r} \Bigl(\prod_{k \neq j} \langle u_{i_k}|v_{i_k}\rangle \nu_{i_j}(\gamma_{u_{i_j},v_{i_j}}) \Bigr),$$
for any $u_{i_j}, v_{i_j} \in V_{i_j}$, $j = 1,\dots,r$, where $\nu_i : \mathfrak{sp}(V_i) \to \mathfrak{sp}(V)^n$ denotes the canonical inclusion into the i^{th}-component, and $\gamma_{u,v}$ has been defined in (2.14).

Example 4.10. The description of $\mathfrak{d}_{2,1}$ (characteristic 3) in (2.18) shows that, with $n = 3$,

$$\mathfrak{d}_{2,1} = V(\emptyset) \oplus V(\{1,2,3\}), \tag{4.11}$$

with even part $V(\emptyset) = \mathfrak{sp}(V_1) \oplus \mathfrak{sp}(V_2) \oplus \mathfrak{sp}(V_3)$, and odd part $V(\{1,2,3\}) = V_1 \otimes V_2 \otimes V_3$, and where

$$[x_\sigma, y_\tau] = \epsilon_{\mathfrak{d}_{2,1}}(\sigma,\tau)\varphi_{\sigma,\tau}(x_\sigma, y_\tau),$$

with

$$\begin{gathered} \epsilon_{\mathfrak{d}_{2,1}}(\emptyset,\emptyset) = \epsilon_{\mathfrak{d}_{2,1}}(\emptyset,\{1,2,3\}) = \epsilon_{\mathfrak{d}_{2,1}}(\{1,2,3\},\emptyset) = 1, \\ \epsilon_{\mathfrak{d}_{2,1}}(\{1,2,3\},\{1,2,3\}) = -1. \end{gathered} \tag{4.12}$$

(Same behavior as $\mathfrak{d}_4$ in [Eldb, (2.18) and (2.19)].)

Take the canonical basic elements $\epsilon_1 = (1,0,0)$, $\epsilon_2 = (0,1,0)$ and $\epsilon_3 = (0,0,1)$ in $\mathbb{Z}^3$, and for any $i = 1,2,3$ consider a symplectic basis $\{v_i, w_i\}$ of

V_i (that is, $\langle v_i|w_i\rangle = 1$), and the basic elements in each $\mathfrak{sp}(V_i)$:

$$\begin{aligned} h_i &= \gamma_{v_i,w_i} : \begin{cases} v_i \mapsto -v_i, \\ w_i \mapsto w_i, \end{cases} \\ e_i &= \gamma_{w_i,w_i} : \begin{cases} v_i \mapsto -2w_i = w_i, \\ w_i \mapsto 0, \end{cases} \\ f_i &= -\gamma_{v_i,v_i} : \begin{cases} v_i \mapsto 0, \\ w_i \mapsto -2v_i = v_i, \end{cases} \end{aligned} \tag{4.13}$$

which satisfy

$$[h_i, e_i] = 2e_i, \quad [h_i, f_i] = -2f_i, \quad [e_i, f_i] = h_i.$$

The Lie superalgebra $\mathfrak{d}_{2,1}$ is $\mathbb{Z}^3$-graded by assigning $\deg w_i = \epsilon_i = -\deg v_i$, $i = 1, 2, 3$. Hence, $\deg h_i = 0$, $\deg e_i = 2\epsilon_i$, $\deg f_i = -2\epsilon_i$, for any i.

Then,

$$\Phi_{\mathfrak{d}_{2,1}} = \{\pm 2\epsilon_i : i = 1, 2, 3\} \cup \{\pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_3\} \tag{4.14}$$

is the set of the nonzero degrees that appear in $\mathfrak{d}_{2,1}$. Moreover, the subalgebra $\mathfrak{h} = kh_1 \oplus kh_2 \oplus kh_3$ is a Cartan subalgebra of $\mathfrak{d}_{2,1}$ and there is a natural homomorphism of abelian groups:

$$\begin{aligned} R : \mathbb{Z}^3 &\longrightarrow \mathfrak{h}^* \\ \epsilon_i &\mapsto R(\epsilon_i)(: h_j \mapsto \delta_{ij}). \end{aligned}$$

The image under R of $\Phi_{\mathfrak{d}_{2,1}}$ is precisely the set of roots of $\mathfrak{h}$ in $\mathfrak{d}_{2,1}$.

Consider the lexicographic order on $\mathbb{Z}^3$ with $\epsilon_1 > \epsilon_2 > \epsilon_3 > 0$. The set of the positive elements in $\Phi_{\mathfrak{d}_{2,1}}$ which are not sums of two positive elements is

$$\Pi_{\mathfrak{d}_{2,1}} = \{\alpha_1 = 2\epsilon_2, \alpha_2 = \epsilon_1 - \epsilon_2 - \epsilon_3, \alpha_3 = 2\epsilon_3\}, \tag{4.15}$$

whose elements are linearly independent over $\mathbb{Z}$ and satisfy that $\Phi_{\mathfrak{d}_{2,1}}$ is contained in $\mathbb{Z}\Pi_{\mathfrak{d}_{2,1}} = \mathbb{Z}\alpha_1 \oplus \mathbb{Z}\alpha_2 \oplus \mathbb{Z}\alpha_3$.

Let Q be the free abelian group $\mathbb{Z}\Pi_{\mathfrak{d}_{2,1}}$, take $\tau = \{2\}$ and consider the following elements in $\mathfrak{d}_{2,1}$:

$$\begin{aligned} &E_1 = e_2, \quad && E_2 = w_1 \otimes v_2 \otimes v_3, \quad && E_3 = e_3, \\ &F_1 = f_2, && F_2 = -v_1 \otimes w_2 \otimes w_3, && F_3 = f_3, \\ &H_1 = h_2, && H_2 = h_1 - h_2 - h_3, && H_3 = h_3. \end{aligned}$$

Then $\big(\mathfrak{h}, \{R(\alpha_1), R(\alpha_2), R(\alpha_3)\}, \{H_1, H_2, H_3\}\big)$ is a realization of the matrix:

$$A_{\mathfrak{d}_{2,1}} = \begin{pmatrix} 2 & -1 & 0 \\ 1 & 0 & 1 \\ 0 & -1 & 2 \end{pmatrix}, \tag{4.16}$$

which is the Cartan matrix D_1 in [Kac77, p. 55], and which corresponds to the Lie superalgebra $D(2,1) \cong D(2,1;1)$, that is, to the orthosymplectic Lie superalgebra $\mathfrak{osp}(4,2)$.

The Lie superalgebra $\mathfrak{d}_{2,1}$ is $\mathbb{Z}$-graded with $(\mathfrak{d}_{2,1})_0 = \mathfrak{h}$, $(\mathfrak{d}_{2,1})_1 = kE_1 \oplus kE_2 \oplus kE_3$, and $(\mathfrak{d}_{2,1})_{-1} = kF_1 \oplus kF_2 \oplus kF_3$. It is easily checked to be simple. Now, Theorem 4.4, with the elements H_i, E_i, F_i above, $i = 1, 2, 3$, gives:

Proposition 4.17. *The Lie superalgebra* $\mathfrak{D}_{2,1}$ *is isomorphic to the contragredient Lie superalgebra* $\mathfrak{g}\bigl(A_{\mathfrak{D}_{2,1}},\{2\}\bigr)$.

5. The Lie superalgebras in the Extended Freudenthal's Magic Square

In [Eldb], most of the Lie algebras in Freudenthal's Magic Square were described, in the split case, by means of the $V(\sigma)$'s and $\varphi_{\sigma,\tau}$'s of the previous section. This is possible too in the superalgebra setting.

This section will be devoted to get such descriptions, or a similar one, for each Lie superalgebra in the Extended Freudenthal's Magic Square, constructed from a couple of split symmetric composition (super)algebras. This will be used to find a description of all these superalgebras as contragredient Lie superalgebras.

Throughout this section, the characteristic of the ground field k will always be assumed to be 3.

5.1. $\mathfrak{g}(S_1,S_{4,2})$**.** The symmetric composition superalgebra $(S_1,\bullet,q)$ is just $k1$, with $1\bullet 1=1$ and $q(1)=1$ (so $b(1,1)=2$). Thus $\mathfrak{tri}(S_1)=0$ and (3.1), (4.11) and (2.25) show that

$$\begin{split}\mathfrak{g}(S_1,S_{4,2})&=\mathfrak{tri}(S_{4,2})\oplus\iota_0(S_1\otimes S_{4,2})\oplus\iota_1(S_1\otimes S_{4,2})\oplus\iota_2(S_1\otimes S_{4,2})\\&=\bigl(V(\emptyset)\oplus V(\{1,2,3\})\bigr)\oplus\bigl(V(\{1,2\})\oplus V(\{3\})\bigr)\\&\qquad\oplus\bigl(V(\{2,3\})\oplus V(\{1\})\bigr)\oplus\bigl(V(\{1,3\})\oplus V(\{2\})\bigr)\\&=\bigoplus_{\sigma\in 2^{\{1,2,3\}}}V(\sigma),\end{split}$$

where $2^{\{1,2,3\}}$ denotes the power set of $\{1,2,3\}$. Besides,

$$\mathfrak{g}(S_1,S_{4,2})_{\bar 0}=\bigoplus_{\substack{\sigma\in 2^{\{1,2,3\}}\\ |\sigma|\text{ even}}}V(\sigma)\qquad\text{and}\qquad\mathfrak{g}(S_1,S_{4,2})_{\bar 1}=\bigoplus_{\substack{\sigma\in 2^{\{1,2,3\}}\\ |\sigma|\text{ odd}}}V(\sigma).$$

By invariance of the Lie bracket under the action of $\mathfrak{sp}(V_1)\oplus\mathfrak{sp}(V_2)\oplus\mathfrak{sp}(V_3)=V(\emptyset)$, it follows that

$$[x_\sigma,y_\tau]=\epsilon(\sigma,\tau)\varphi_{\sigma,\tau}(x_\sigma,y_\tau),\tag{5.1}$$

for any $\sigma,\tau\in 2^{\{1,2,3\}}$, for a suitable map

$$\epsilon:2^{\{1,2,3\}}\times 2^{\{1,2,3,\}}\to k^\times.$$

The multiplication $\iota_i(1\otimes S_{4,2})\times\iota_{i+1}(1\otimes S_{4,2})\to\iota_{i+2}(1\otimes S_{4,2})$ is given in (2.26), and for any $x,y\in S_{4,2}$ (see (3.2)):

$$[\iota_0(1\otimes x),\iota_0(1\otimes y)]=2t_{x,y}=\bigl(2\sigma_{x,y},b(x,y)1-2r_xl_y,b(x,y)1-2l_xr_y\bigr),$$

as $b(1,1)=2$ in S_1. This element is identified with the element $2\rho^{-1}(\sigma_{x,y})$ in (2.19), which is given by (2.20). By cyclic symmetry, one completes the information about the map ϵ in (5.1), displayed in Table 3, which is exactly Table 2 in [Eldb], corresponding to $\mathfrak{g}(S_1,S_8)$, but with the index 4 taken out. From this description it readily follows that $\mathfrak{g}(S_1,S_{4,2})$ is simple.

ϵ	$\emptyset$	$\{1,2\}$	$\{2,3\}$	$\{1,3\}$	$\{1,2,3\}$	$\{3\}$	$\{1\}$	$\{2\}$
$\emptyset$	1	1	1	1	1	1	1	1
$\{1,2\}$	1	-2	1	1	1	-2	-1	-1
$\{2,3\}$	1	1	-2	1	1	-1	-2	-1
$\{1,3\}$	1	1	1	-2	1	-1	-1	-2
$\{1,2,3\}$	1	-1	-1	-1	-1	1	1	1
$\{3\}$	1	2	-1	-1	-1	-2	-1	-1
$\{1\}$	1	-1	2	-1	-1	-1	-2	-1
$\{2\}$	1	-1	-1	2	-1	-1	-1	-2

TABLE 3. $\epsilon_{S_1,S_{4,2}}$

In general, in order to obtain a description of the Lie superalgebra $\mathfrak{g}(S,S')$ in the Extended Freudenthal's Magic Square as contragredient Lie superalgebras, the complete description of the map ϵ in (5.1) is not necessary and will not be given. A detailed description of these maps appear in [Cun06].

Now, with the notations and conventions in Example 4.10, the Lie superalgebra $\mathfrak{g} = \mathfrak{g}(S_1, S_{4,2})$ is $\mathbb{Z}^3$-graded, and the set of nonzero degrees that appear is

$$\Phi_{S_1,S_{4,2}} = \bigl\{\pm 2\epsilon_i, \pm\epsilon_i \pm \epsilon_j, \pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_3, \pm\epsilon_i : 1 \leq i < j \leq 3\bigr\}.$$

The subalgebra $\mathfrak{h} = kh_1 \oplus kh_2 \oplus kh_3$ (recall (4.13) that $h_i = \gamma_{v_i,w_i}$ belongs to $\mathfrak{sp}(V_i) \subseteq V(\emptyset)$, where $\{v_i, w_i\}$ is a fixed symplectic basis of V_i) is a Cartan subalgebra of $\mathfrak{g}$ and the image of $\Phi_{S_1,S_{4,2}}$ under the natural homomorphism of abelian groups

$$\begin{aligned} R : \mathbb{Z}^3 &\longrightarrow \mathfrak{h}^* \\ \epsilon_i &\mapsto R(\epsilon_i)(: h_j \mapsto \delta_{ij}) \end{aligned}$$

is precisely the set of roots of $\mathfrak{g}$ relative to $\mathfrak{h}$.

Consider the lexicographic order on $\mathbb{Z}^3$ with $\epsilon_1 > \epsilon_2 > \epsilon_3 > 0$. The set of positive elements in $\Phi_{S_1,S_{4,2}}$ which are not sums of positive elements (the *irreducible* degrees) is

$$\Pi = \{\alpha_1 = \epsilon_2 - \epsilon_3, \alpha_2 = \epsilon_3, \alpha_3 = \epsilon_1 - \epsilon_2 - \epsilon_3\},$$

which is a $\mathbb{Z}$-basis of $\mathbb{Z}^3$.

Then $\bigl(\mathfrak{h}, \{R(\alpha_1), R(\alpha_2), R(\alpha_3)\}, \{h_2 - h_3, 2h_3, -h_1 + h_2 + h_3\}\bigr)$ is a realization of the matrix

$$A_{S_1,S_{4,2}} = \begin{pmatrix} 2 & -1 & 0 \\ 1 & 2 & 1 \\ 0 & 1 & 0 \end{pmatrix},$$

with associated Dynkin diagram (using the conventions in [Kac77, Tables IV and V])



With $\tau = \{2,3\}$, consider the generators

$$\begin{aligned} &H_1 = h_2 - h_3, \quad H_2 = 2h_3, \quad H_3 = -h_1 + h_2 + h_3,\\ &E_1 = w_2 \otimes v_3, \quad E_2 = w_3, \quad E_3 = w_1 \otimes v_2 \otimes v_3,\\ &F_1 = v_2 \otimes w_3, \quad F_2 = -v_3, \quad F_3 = -v_1 \otimes w_2 \otimes w_3. \end{aligned}$$

As for Proposition 4.17, Theorem 4.4 implies the following description of $\mathfrak{g}(S_1, S_{4,2})$:

Proposition 5.2. *The Lie superalgebra $\mathfrak{g}(S_1, S_{4,2})$ is isomorphic to the contragredient Lie superalgebra $\mathfrak{g}\big(A_{S_1,S_{4,2}}, \{2,3\}\big)$.*

Moreover, the set of even and odd nonzero degrees are

$$\begin{aligned} \big(\Phi_{S_1,S_{4,2}}\big)_{\bar 0} &= \{\pm 2\epsilon_i, \pm\epsilon_i \pm \epsilon_j : 1 \le i < j \le 3\},\\ \big(\Phi_{S_1,S_{4,2}}\big)_{\bar 1} &= \{\pm\epsilon_i, \pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_3 : 1 \le i \le 3\}. \end{aligned}$$

The set of "irreducible"even degrees for the lexicographic order considered is

$$\Pi_{\bar 0} = \{\beta_1 = \epsilon_1 - \epsilon_2, \beta_2 = \epsilon_2 - \epsilon_3, \beta_3 = 2\epsilon_3\},$$

and it can be concluded from here that the even part $\mathfrak{g}_{\bar 0} = \mathfrak{g}(S_1, S_{4,2})_{\bar 0}$ is isomorphic to the contragredient Lie algebra with Cartan matrix

$$\begin{pmatrix} 2 & -1 & 0\\ -1 & 2 & -2\\ 0 & -1 & 2 \end{pmatrix}.$$

That is, $\mathfrak{g}(S_1, S_{4,2})_{\bar 0}$ is isomorphic to the symplectic Lie algebra $\mathfrak{sp}_6$ (type C_3). Here the appropriate basis of $\mathfrak{h}$ is

$$\{\tilde H_1 = h_1 - h_2, \tilde H_2 = h_2 - h_3, \tilde H_3 = h_3\}.$$

Also, the set of simple roots $R(\Pi_{\bar 0})$ of $\mathfrak{g}_{\bar 0}$ induces a triangular decomposition $\mathfrak{g}_{\bar 0} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+$, where $\mathfrak{n}^+ = \oplus_{0<\alpha\in(\Phi_{S_1,S_{4,2}})_{\bar 0}} (\mathfrak{g}_{\bar 0})_{R(\alpha)}$ is the sum of root spaces (nonzero homogeneous components in the $\mathbb{Z}^3$-grading) corresponding to the positive degrees in $(\Phi_{S_1,S_{4,2}})_{\bar 0}$, and similarly for $\mathfrak{n}^-$. The $\mathfrak{g}_{\bar 0}$-module $\mathfrak{g}_{\bar 1}$ is $\mathbb{Z}^3$-graded consistently with the action of $\mathfrak{g}_{\bar 0}$, it is easily seen to be irreducible, and to contain the highest weight vector $w_1 \otimes w_2 \otimes w_3$, that is, $[\mathfrak{n}^+, w_1 \otimes w_2 \otimes w_3] = 0$. Its highest weight is $R(\epsilon_1 + \epsilon_2 + \epsilon_3) = \omega_3$, which satisfies $\omega_3(\tilde H_1) = 0 = \omega_3(\tilde H_2)$, $\omega_3(\tilde H_3) = 1$.

Since the proof of the uniqueness result in [Hum72, Theorem A in §20.3] remains valid in this setting (note that instead of grading over the $\mathbb{Z}$-linear combinations of weights, which are elements of $\mathfrak{h}^*$, we grade over a true lattice $\mathbb{Z}^3$), $\mathfrak{g}_{\bar 1}$ is the unique $\mathfrak{g}_{\bar 0}$-module with such a highest weight. Denote it by $V(\omega_3)$. Note that $\dim \mathfrak{g}_{\bar 1} = 14$.

Let us give the most natural presentation of this module. Consider the matrix Lie algebra $\mathfrak{sp}_6$, and its natural six-dimensional module, which is endowed with an alternating invariant bilinear form $\{.|.\}$. Let $\{a_1, a_2, a_3, b_1, b_2, b_3\}$ be a symplectic basis (that is, $\{a_i|b_j\} = \delta_{ij}$, $\{a_i|a_j\} = 0 = \{b_i|b_j\}$ for any i, j). With $\gamma_{x,y} = \{x|.\}y + \{y|.\}x$, the subspace spanned by γ_{a_i,b_i}, $i = 1, 2, 3$ is a Cartan subalgebra of $\mathfrak{sp}(W) \simeq \mathfrak{sp}_6$ and the weights of W are $\{\pm\delta_i : i = 1, 2, 3\}$, where $\delta_i(\gamma_{a_j,b_j}) = \delta_{ij}$ for any i, j.

Consider the $\mathfrak{sp}_6$-module $\bigwedge^3 W$ and the homomorphism

$$\begin{aligned}\varphi : \bigwedge^3 W &\longrightarrow W\\ z_1 \wedge z_2 \wedge z_3 &\mapsto \{z_1|z_2\}z_3 + \{z_2|z_3\}z_1 + \{z_3|z_1\}z_2.\end{aligned}$$

Then $\dim\ker\varphi = 14$, the weights of $\ker\varphi$ are $\pm\delta_1 \pm \delta_2 \pm \delta_3$ and $\pm\delta_i$ ($i = 1,2,3$), all of them of multiplicity 1. The element $b_1 \wedge b_2 \wedge b_3$ is a highest weight vector of weight $\delta_1 + \delta_2 + \delta_3 = \omega_3$ and $\ker\varphi$ is irreducible. Therefore, up to isomorphism $V(\omega_3) = \ker\varphi$.

Let us summarize the above discussion:

Proposition 5.3. *The Lie superalgebra $\mathfrak{g}(S_1, S_{4,2})$ is simple with even part isomorphic to the symplectic Lie algebra $\mathfrak{sp}_6$ and odd part isomorphic to the irreducible module of dimension 14 above.*

There is no counterpart in characteristic 0 (see [Kac77]) to this simple Lie superalgebra.

5.2. $\mathfrak{g}(S_4, S_{4,2})$.

The symmetric composition superalgebra S_4 is the even part of $S_{4,2}$, and its triality Lie algebra is given too by the even part of the triality Lie superalgebra of $S_{4,2}$. Therefore, six copies of V are needed: V_1, V_2 and V_3 for S_4, and V_4, V_5 and V_6 for $S_{4,2}$.

With the same sort of arguments used so far, one has:

$$\begin{aligned}\mathfrak{g}(S_4, S_{4,2}) &= \bigl(\mathfrak{tri}(S_4) \oplus \mathfrak{tri}(S_{4,2})\bigr) \oplus \bigl(\oplus_{i=0}^2 \iota_i(S_4 \otimes S_{4,2})\bigr)\\ &= \oplus_{\sigma\in\mathcal{S}_{S_4,S_{4,2}}} V(\sigma),\end{aligned}$$

where

$$\begin{aligned}\mathcal{S} = \mathcal{S}_{S_4,S_{4,2}} = \bigl\{&\emptyset, \{4,5,6\},\\ &\{1,2,4,5\}, \{1,2,6\}, \{2,3,5,6\}, \{2,3,4\}, \{1,3,4,6\}, \{1,3,5\}\bigr\}.\end{aligned}$$

For instance,

$$\iota_1(S_4 \otimes S_{4,2}) = (V_2 \otimes V_3) \otimes \bigl((V_5 \otimes V_6) \oplus V_4\bigr) = V\bigl(\{2,3,5,6\}\bigr) \oplus V\bigl(\{2,3,4\}\bigr).$$

Like in all the other cases, the even (respectively odd) part is the sum of the $V(\sigma)$'s with σ containing an even (resp. odd) number of elements.

The multiplication presents the form in (5.1) for a suitable map $\epsilon : \mathcal{S}\times\mathcal{S} \to k^\times$. Here the nonzero even and odd degrees are:

$$\begin{aligned}\Phi_{\bar 0} = \{&\pm 2\epsilon_i : 1 \le i \le 6\}\\ &\cup \{\pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_4 \pm \epsilon_5, \pm\epsilon_2 \pm \epsilon_3 \pm \epsilon_5 \pm \epsilon_6, \pm\epsilon_1 \pm \epsilon_3 \pm \epsilon_4 \pm \epsilon_6\},\end{aligned}$$

$$\Phi_{\bar 1} = \{\pm\epsilon_4 \pm \epsilon_5 \pm \epsilon_6, \pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_6, \pm\epsilon_2 \pm \epsilon_3 \pm \epsilon_4, \pm\epsilon_1 \pm \epsilon_3 \pm \epsilon_5\}.$$

With the lexicographic order given by $0 < \epsilon_1 < \cdots < \epsilon_6$, the set of irreducible degrees is

$$\begin{aligned}\Pi = \{&\alpha_1 = 2\epsilon_1, \alpha_2 = \epsilon_5 - \epsilon_4 - \epsilon_2 - \epsilon_1, \alpha_3 = 2\epsilon_2,\\ &\alpha_4 = \epsilon_4 - \epsilon_3 - \epsilon_2, \alpha_5 = 2\epsilon_3, \alpha_6 = \epsilon_6 - \epsilon_5 - \epsilon_4\},\end{aligned}$$

which is a linearly independent set over $\mathbb{Z}$. Then Φ is contained in $Q = \mathbb{Z}\Pi$ (which is isomorphic to $\mathbb{Z}^6$), so that any positive element in Φ is a sum of elements in Π. Consider the elements (same conventions as before)

$$H_1 = h_1,\ H_2 = -h_5 + h_4 + h_2 + h_1,\ H_3 = h_2,\ H_4 = h_4 - h_3 - h_2,\\ H_5 = h_3,\ H_6 = h_6 - h_5 - h_4,$$

which give a realization

$$\bigl(\mathfrak{h}, \{R(\alpha_1), R(\alpha_2), R(\alpha_3), R(\alpha_4), R(\alpha_5), R(\alpha_6)\}, \{H_1, H_2, H_3, H_4, H_5, H_6\}\bigr)$$

of the regular matrix (with associated Dynkin diagram):

$$A_{S_4,S_{4,2}} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & -1 \\ 0 & 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix},$$



Then:

Proposition 5.4. *The Lie superalgebra* $\mathfrak{g}(S_4, S_{4,2})$ *is isomorphic to the contragredient Lie superalgebra* $\mathfrak{g}\bigl(A_{S_4,S_{4,2}}, \{4,6\}\bigr)$.

Also, the set of irreducible even degrees is:

$$\Pi_{\bar 0} = \{\beta_1 = 2\epsilon_3, \beta_2 = \epsilon_6 - \epsilon_5 - \epsilon_3 - \epsilon_2, \beta_3 = 2\epsilon_2,\\ \beta_4 = \epsilon_5 - \epsilon_4 - \epsilon_2 - \epsilon_1, \beta_5 = 2\epsilon_1, \beta_6 = 2\epsilon_4\},$$

with associated Cartan matrix of type D_6. The odd part is an irreducible module for the even part, with highest weight vector $R(\epsilon_4 + \epsilon_5 + \epsilon_6) = \omega_6$, and the same sort of arguments used for $\mathfrak{g}(S_1, S_{4,2})$ gives:

Proposition 5.5. *The Lie superalgebra* $\mathfrak{g}(S_4, S_{4,2})$ *is simple with even part isomorphic to the orthogonal Lie algebra* $\mathfrak{so}_{12}$*, and odd part isomorphic to the spin module for* $\mathfrak{so}_{12}$.

Note that the even part is just $\mathfrak{g}(S_4, S_4)$, which was determined in [Eldb].

Corollary 5.6. *The Lie superalgebra* $\mathfrak{g}(S_4, S_{4,2})$ *is isomorphic to the Lie superalgebra in* [Eld06b, theorem 3.2(v)] *and* [Elda, Theorem 4.1(ii) $(l = 6)$].

5.3. $\mathfrak{g}(S_8, S_{4,2})$. Here, four copies of V are needed for S_8 (see [Eldb, (2.14)]) and three copies for $S_{4,2}$. The indices 1, 2, 3 and 4 will be used for S_8, while 5, 6 and 7 will be reserved for $S_{4,2}$. Then

$$\begin{aligned}\mathfrak{g}(S_8, S_{4,2}) &= \bigl(\mathfrak{tri}(S_8) \oplus \mathfrak{tri}(S_{4,2})\bigr) \oplus \bigl(\oplus_{i=0}^{2} \iota_i(S_8 \otimes S_{4,2})\bigr)\\ &= \oplus_{\sigma \in \mathcal{S}_{S_8,S_{4,2}}} V(\sigma),\end{aligned}$$

where

$$\begin{aligned}\mathcal{S} = \mathcal{S}_{S_8,S_{4,2}} = \bigl\{&\emptyset, \{1,2,3,4\}, \{5,6,7\},\\ &\{1,2,5,6\}, \{3,4,5,6\}, \{1,2,7\}, \{3,4,7\},\\ &\{2,3,6,7\}, \{1,4,6,7\}, \{2,3,5\}, \{1,4,5\},\\ &\{1,3,5,7\}, \{2,4,5,7\}, \{1,3,6\}, \{2,4,6\}\bigr\}.\end{aligned}$$

Hence,

$$\begin{aligned}\Phi_{\bar 0} = \bigl\{&\pm 2\epsilon_i : i = 1, \dots, 7\bigr\} \cup \bigl\{\pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_3 \pm \epsilon_4,\\ &\pm \epsilon_1 \pm \epsilon_2 \pm \epsilon_5 \pm \epsilon_6, \pm\epsilon_3 \pm \epsilon_4 \pm \epsilon_5 \pm \epsilon_6, \pm\epsilon_2 \pm \epsilon_3 \pm \epsilon_6 \pm \epsilon_7,\\ &\pm \epsilon_1 \pm \epsilon_4 \pm \epsilon_6 \pm \epsilon_7, \pm\epsilon_1 \pm \epsilon_3 \pm \epsilon_5 \pm \epsilon_7, \pm\epsilon_2 \pm \epsilon_4 \pm \epsilon_5 \pm \epsilon_7\bigr\},\end{aligned}$$

$$\begin{aligned}\Phi_{\bar 1} = \bigl\{&\pm\epsilon_5 \pm \epsilon_6 \pm \epsilon_7, \pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_7, \pm\epsilon_3 \pm \epsilon_4 \pm \epsilon_7, \pm\epsilon_2 \pm \epsilon_3 \pm \epsilon_5,\\ &\pm \epsilon_1 \pm \epsilon_4 \pm \epsilon_5, \pm\epsilon_1 \pm \epsilon_3 \pm \epsilon_6, \pm\epsilon_2 \pm \epsilon_4 \pm \epsilon_6\bigr\},\end{aligned}$$

and with the lexicographic order with $0 < \epsilon_1 < \cdots < \epsilon_7$:

$$\begin{aligned}\Pi = \bigl\{&\alpha_1 = \epsilon_6 - \epsilon_5 - \epsilon_4 - \epsilon_3, \alpha_2 = 2\epsilon_2, \alpha_3 = 2\epsilon_3,\\ &\alpha_4 = \epsilon_4 - \epsilon_3 - \epsilon_2 - \epsilon_1, \alpha_5 = 2\epsilon_1, \alpha_6 = \epsilon_5 - \epsilon_4 - \epsilon_1, \alpha_7 = \epsilon_7 - \epsilon_6 - \epsilon_5\bigr\},\end{aligned}$$

which is a $\mathbb{Z}$-linearly independent set with $\Phi \subseteq \mathbb{Z}\Pi$, so that any positive element in Φ is a sum of elements in Π. The associated matrix and Dynkin diagram are here:

$$A_{S_8,S_{4,2}} = \begin{pmatrix} 2 & 0 & -1 & 0 & 0 & 0 & 0\\ 0 & 2 & 0 & -1 & 0 & 0 & 0\\ -1 & 0 & 2 & -1 & 0 & 0 & 0\\ 0 & -1 & -1 & 2 & -1 & 0 & 0\\ 0 & 0 & 0 & -1 & 2 & -1 & 0\\ 0 & 0 & 0 & 0 & 1 & 0 & -1\\ 0 & 0 & 0 & 0 & 0 & -1 & 0 \end{pmatrix},$$



Proposition 5.7. *The Lie superalgebra* $\mathfrak{g}(S_8, S_{4,2})$ *is isomorphic to the contragredient Lie superalgebra* $\mathfrak{g}\bigl(A_{S_8,S_{4,2}}, \{6, 7\}\bigr)$.

Also, in this situation:

$$\begin{aligned}\Pi_{\bar 0} = \bigl\{&\beta_1 = \epsilon_7 - \epsilon_6 - \epsilon_4 - \epsilon_1, \beta_2 = 2\epsilon_2, \beta_3 = 2\epsilon_1,\\ &\beta_4 = \epsilon_4 - \epsilon_3 - \epsilon_2 - \epsilon_1, \beta_5 = 2\epsilon_3, \beta_6 = \epsilon_6 - \epsilon_5 - \epsilon_4 - \epsilon_3, \beta_7 = 2\epsilon_5\bigr\},\end{aligned}$$

whose associated Cartan matrix is of type E_7 (this also follows since $\mathfrak{g}(S_8, S_{4,2})_{\bar 0}$ is isomorphic to $\mathfrak{g}(S_8, S_4)$, which was computed in [Eldb]).

The odd part, which has dimension $7 \times 8 = 56$, is an irreducible module with highest weight $R(\epsilon_7 + \epsilon_6 + \epsilon_5) = \omega_7$. Thus,

Proposition 5.8. $\mathfrak{g}(S_8, S_{4,2})$ *is a simple Lie superalgebra, whose even part is isomorphic to the split simple Lie algebra of type* E_7*, and whose odd part is its* 56*-dimensional irreducible module* $V(\omega_7)$.

5.4. $\mathfrak{g}(S_{4,2}, S_{4,2})$.

For $\mathfrak{g}(S_{4,2}, S_{4,2})$, the indices 1, 2 and 3 will refer to the copies of V associated to the first copy of $S_{4,2}$, while the indices 4, 5 and 6 will refer to the copies of V related to the second copy of $S_{4,2}$. Then:

$$\begin{aligned}\mathfrak{g}(S_{4,2}, S_{4,2}) &= \bigl(\mathfrak{tri}(S_{4,2}) \oplus \mathfrak{tri}(S_{4,2})\bigr) \oplus \bigl(\oplus_{i=0}^2 \iota_i(S_{4,2} \otimes S_{4,2})\bigr)\\ &= \oplus_{\sigma\in\mathcal{S}_{S_{4,2},S_{4,2}}} V(\sigma),\end{aligned}$$

where

$$\begin{aligned}\mathcal{S}_{S_{4,2},S_{4,2}} = \big\{&\emptyset,\{1,2,3\},\{4,5,6\},\\ &\{1,2,4,5\},\{3,6\},\{1,2,6\},\{3,4,5\},\\ &\{2,3,5,6,\},\{1,4\},\{2,3,4\},\{1,5,6\},\\ &\{1,3,4,6\},\{2,5\},\{1,3,5\},\{2,4,6\}\big\}.\end{aligned}$$

Hence,

$$\begin{aligned}\Phi_{\bar 0} = \big\{&\pm 2\epsilon_i : 1\le i\le 6\big\}\cup\big\{\pm\epsilon_1\pm\epsilon_2\pm\epsilon_4\pm\epsilon_5, \pm\epsilon_2\pm\epsilon_3\pm\epsilon_5\pm\epsilon_6,\\ &\pm\epsilon_1\pm\epsilon_3\pm\epsilon_4\pm\epsilon_6, \pm\epsilon_3\pm\epsilon_6, \pm\epsilon_1\pm\epsilon_4, \pm\epsilon_2\pm\epsilon_5\big\},\end{aligned}$$

$$\begin{aligned}\Phi_{\bar 1} = \big\{&\pm\epsilon_1\pm\epsilon_2\pm\epsilon_3, \pm\epsilon_4\pm\epsilon_5\pm\epsilon_6, \pm\epsilon_1\pm\epsilon_2\pm\epsilon_6, \pm\epsilon_3\pm\epsilon_4\pm\epsilon_5,\\ &\pm\epsilon_2\pm\epsilon_3\pm\epsilon_4, \pm\epsilon_1\pm\epsilon_5\pm\epsilon_6, \pm\epsilon_1\pm\epsilon_3\pm\epsilon_5, \pm\epsilon_2\pm\epsilon_4\pm\epsilon_6\big\}.\end{aligned}$$

And with the lexicographic order with $\epsilon_1 > \cdots > \epsilon_6 > 0$,

$$\begin{aligned}\Pi = \big\{&\alpha_1 = \epsilon_1-\epsilon_2-\epsilon_3, \alpha_2 = \epsilon_3-\epsilon_4-\epsilon_5, \alpha_3 = 2\epsilon_5,\\ &\alpha_4 = \epsilon_4-\epsilon_5-\epsilon_6, \alpha_5 = 2\epsilon_6, \alpha_6 = \epsilon_2-\epsilon_3-\epsilon_4\big\},\end{aligned}$$

which is a linearly independent set with $\Phi \subseteq \mathbb{Z}\Pi$, so that any positive element in Φ is a sum of elements in Π. The associated matrix and Dynkin diagram are:

$$A_{S_{4,2},S_{4,2}} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0\\ 1 & 0 & -1 & 0 & 0 & 0\\ 0 & -1 & 2 & -1 & 0 & 0\\ 0 & 0 & -1 & 0 & -1 & -1\\ 0 & 0 & 0 & -1 & 2 & 0\\ 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix},$$



Proposition 5.9. *The Lie superalgebra* $\mathfrak{g}(S_{4,2},S_{4,2})$ *is isomorphic to the contragredient Lie superalgebra* $\mathfrak{g}\big(A_{S_{4,2},S_{4,2}},\{1,2,4,6\}\big)$.

Also,

$$\begin{aligned}\Pi_{\bar 0} = \big\{&\beta_1 = 2\epsilon_4, \beta_2 = \epsilon_1-\epsilon_2-\epsilon_4-\epsilon_5, \beta_3 = 2\epsilon_5,\\ &\beta_4 = \epsilon_2-\epsilon_3-\epsilon_5-\epsilon_6, \beta_5 = 2\epsilon_6, \beta_6 = \epsilon_3-\epsilon_6\big\},\end{aligned}$$

with associated Cartan matrix of type B_6. The odd part is an irreducible module for the even part with highest weight $R(\epsilon_1+\epsilon_2+\epsilon_3) = \omega_6$, so by uniqueness, it is the spin module for the even part:

Proposition 5.10. $\mathfrak{g}(S_{4,2},S_{4,2})$ *is a simple Lie superalgebra whose even part is isomorphic to the orthogonal Lie algebra* $\mathfrak{so}_{13}$ *and whose odd part is the spin module for its even part.*

Corollary 5.11. *The Lie superalgebra* $\mathfrak{g}(S_{4,2},S_{4,2})$ *is isomorphic to the Lie superalgebra in* [Elda, Theorem 3.1(ii) $(l = 6)$].

5.5. **$\mathfrak{g}(S_{1,2},S_{4,2})$.** Here there is just one copy of V involved in $S_{1,2}$, which will carry index 1, and three copies, with indices 2, 3 and 4, in $S_{4,2}$. Also, the triality Lie superalgebra $\mathfrak{tri}(S_{1,2})$ will be identified to the superalgebra $\mathfrak{b}_{0,1}$ in (2.17). Then

$$\begin{aligned}\mathfrak{g}(S_{1,2},S_{4,2}) &= \bigl(\mathfrak{tri}(S_{1,2})\oplus\mathfrak{tri}(S_{4,2})\bigr)\oplus\bigl(\oplus_{i=0}^{2}\iota_i(S_{1,2}\otimes S_{4,2})\bigr)\\ &= \oplus_{\sigma\in\mathcal{S}_{S_{1,2},S_{4,2}}}V(\sigma),\end{aligned}$$

with

$$\begin{aligned}\mathcal{S}_{S_{1,2},S_{4,2}} = \bigl\{&\emptyset,\{1\},\{2,3,4\},\{1,4\},\{1,2,3\},\{4\},\\ &\{3,4\},\{1,2\},\{1,3,4\},\{2\},\{2,4\},\{1,3\},\{1,2,4\},\{3\}\bigr\},\end{aligned}$$

and

$$\begin{aligned}\Phi_{\bar 0} &= \{\pm 2\epsilon_i : 1\le i\le 4\}\cup\{\pm\epsilon_i\pm\epsilon_j : 1\le i<j\le 4\},\\ \Phi_{\bar 1} &= \{\pm\epsilon_i\pm\epsilon_j\pm\epsilon_k : 1\le i<j<k\le 4\}\cup\{\pm\epsilon_i : 1\le i\le 4\}.\end{aligned}$$

With the lexicographic order $\epsilon_1>\epsilon_2>\epsilon_3>\epsilon_4>0$,

$$\Pi=\{\alpha_1=\epsilon_1-\epsilon_2-\epsilon_3,\alpha_2=\epsilon_3-\epsilon_4,\alpha_3=\epsilon_4,\alpha_4=\epsilon_2-\epsilon_3-\epsilon_4\},$$

which is a linearly independent set with $\Phi\subseteq\mathbb{Z}\Pi$, so that any positive element in Φ is a sum of elements in Π. The associated matrix and Dynkin diagram are:

$$A_{S_{1,2},S_{4,2}}=\begin{pmatrix}0&1&0&0\\-1&2&-1&0\\0&-2&2&-2\\0&0&1&0\end{pmatrix},$$



and, therefore:

Proposition 5.12. *The Lie superalgebra* $\mathfrak{g}(S_{1,2},S_{4,2})$ *is isomorphic to the contragredient Lie superalgebra* $\mathfrak{g}\bigl(A_{S_{1,2},S_{4,2}},\{1,3,4\}\bigr)$.

Also,

$$\Pi_{\bar 0}=\{\beta_1=\epsilon_1-\epsilon_2,\beta_2=\epsilon_2-\epsilon_3,\beta_3=\epsilon_3-\epsilon_4,\beta_4=2\epsilon_4\},$$

with associated Cartan matrix of type C_4. The odd part is an irreducible module of dimension $4\times 8+4\times 2=40$, with highest weight $R(\epsilon_1+\epsilon_2+\epsilon_3)=\omega_3$.

As for the Lie superalgebra $\mathfrak{g}(S_1,S_{4,2})$, the irreducible module for the symplectic Lie algebra $\mathfrak{sp}_8$ with this highest weight is obtained as follows. Let W be the natural eight dimensional module for $\mathfrak{sp}_8$, and let $\varphi:\bigwedge^3 W\to W$ be the linear map such that $\varphi(z_1\wedge z_2\wedge z_3)=\{z_1|z_2\}z_3+\{z_2|z_3\}z_1+\{z_3|z_1\}z_2$. This time $\ker\varphi$ is not irreducible but, since the characteristic is 3, contains the irreducible submodule $\tilde W=\{\sum_{i=1}^4 a_i\wedge b_i\wedge z_i : z\in W\}$, which is isomorphic to W (here $\{a_i,b_i : i=1,\dots,4\}$ is a symplectic basis of W). Then $\ker\varphi/\tilde W$ is an irreducible module of dimension 40 and the class of $b_1\wedge b_2\wedge b_3$ modulo $\tilde W$ is a highest weight vector of weight ω_3.

Proposition 5.13. $\mathfrak{g}(S_{1,2},S_{4,2})$ *is a simple Lie superalgebra with even part isomorphic to the symplectic Lie algebra* $\mathfrak{sp}_8$ *and odd part isomorphic to the irreducible module of dimension* 40 *above.*

5.6. **$\mathfrak{g}(S_2, S_{4,2})$.** The symmetric composition superalgebra S_2 has a basis $\{e^+, e^-\}$ with multiplication given by

$$e^\pm \bullet e^\pm = e^\mp, \qquad e^\pm \bullet e^\mp = 0,$$

with norm given by $q(e^\pm) = 0$, and $b(e^+, e^-) = 1$. The orthogonal Lie algebra $\mathfrak{so}(S_2, q)$ is spanned by $\phi = \sigma_{e^-,e^+}$, which satisfies $\phi(e^\pm) = \pm e^\pm$, and its triality Lie algebra is (see [Eldb, 3.4]):

$$\mathfrak{tri}(S_2) = \{(\mu_0\phi, \mu_1\phi, \mu_2\phi) : \mu_0, \mu_1, \mu_2 \in k,\ \mu_0 + \mu_1 + \mu_2 = 0\}.$$

Besides,

$$\begin{aligned} t_{e^-,e^+} &= \bigl(\sigma_{e^-,e^+}, \frac{1}{2}b(e^-, e^+)1 - r_{e^-}l_{e^+}, \frac{1}{2}b(e^-, e^+)1 - l_{e^-}r_{e^+}\bigr) \\ &= (\phi, \phi, \phi), \end{aligned}$$

(since the characteristic is 3). Consider the two dimensional abelian Lie algebra

$$\mathfrak{t} = \{(\mu_0, \mu_1, \mu_2) \in k^3 : \mu_0 + \mu_1 + \mu_2 = 0\},$$

with basis $\{t_1 = (1, -1, 0), t_2 = (0, 1, -1)\}$. Then $\mathfrak{tri}(S_2)$ is isomorphic to $\mathfrak{t}$ and the action of $\mathfrak{tri}(S_2)$ on each $\iota_i(S_2)$ becomes:

$$(\mu_0, \mu_1, \mu_2).\iota_i(e^\pm) = \pm\mu_i\iota_i(e^\pm).$$

Here,

$$\mathfrak{g} = \mathfrak{g}(S_2, S_{4,2}) = \bigl(\mathfrak{tri}(S_2) \oplus \mathfrak{tri}(S_{4,2})\bigr) \oplus \bigl(\oplus_{i=0}^2 \iota_i(S_2 \otimes S_{4,2})\bigr),$$

so, with standard identifications,

$$\begin{aligned} \mathfrak{g}_{\bar 0} &= \bigl(\mathfrak{t} \oplus \mathfrak{sp}(V_1) \oplus \mathfrak{sp}(V_2) \oplus \mathfrak{sp}(V_3)\bigr) \oplus \\ &\qquad \bigl(S_2^0 \otimes V_1 \otimes V_2\bigr) \oplus \bigl(S_2^1 \otimes V_2 \otimes V_3\bigr) \oplus \bigl(S_2^2 \otimes V_1 \otimes V_3\bigr) \\ \mathfrak{g}_{\bar 1} &= \bigl(V_1 \otimes V_2 \otimes V_3\bigr) \oplus \bigl(S_2^0 \otimes V_3\bigr) \oplus \bigl(S_2^1 \otimes V_1\bigr) \oplus \bigl(S_2^2 \otimes V_2\bigr), \end{aligned}$$

where, for any $i = 0, 1, 2$, S_2^i is just a copy of S_2, with basic elements $e_i^\pm$.

Consider the free $\mathbb{Z}$-module F with basis $\{\delta_1, \delta_2, \epsilon_1, \epsilon_2, \epsilon_3\}$. The Lie superalgebra $\mathfrak{g}$ is graded over F ($\cong \mathbb{Z}^5$) by assigning degree ϵ_i to w_i, $-\epsilon_i$ to v_i, $i = 1, 2, 3$, as in Example 4.10, and

$$\deg(e_0^\pm) = \pm\delta_1, \quad \deg(e_1^\pm) = \pm\delta_2, \quad \deg(e_2^\pm) = \mp(\delta_1 + \delta_2). \tag{5.14}$$

In this way, the sets of even and odd nonzero degrees that appear in $\mathfrak{g}(S_2, S_{4,2})$ are:

$$\begin{aligned} \Phi_{\bar 0} &= \{\pm 2\epsilon_i : i = 1, 2, 3\} \\ &\qquad \cup \{\pm\delta_1 \pm \epsilon_1 \pm \epsilon_2, \pm\delta_2 \pm \epsilon_2 \pm \epsilon_3, \pm(\delta_1 + \delta_2) \pm \epsilon_1 \pm \epsilon_3\}, \end{aligned}$$

$$\Phi_{\bar 1} = \{\pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_3, \pm\delta_1 \pm \epsilon_3, \pm\delta_2 \pm \epsilon_1, \pm(\delta_1 + \delta_2) \pm \epsilon_2\}.$$

The abelian subalgebra $\mathfrak{h} = kt_1 \oplus kt_2 \oplus kh_1 \oplus kh_2 \oplus kh_3$ ($h_i = \gamma_{v_i,w_i}$, $i = 1, 2, 3$), is a Cartan subalgebra of $\mathfrak{g}(S_2, S_{4,2})$ and the image of $\Phi_{\bar 0} \cup \Phi_{\bar 1}$ under the natural homomorphism of abelian groups

$$\begin{aligned} R : F &\longrightarrow \mathfrak{h}^* \\ \delta_1 &\mapsto R(\delta_1)\,(: t_1 \mapsto 1,\ t_2 \mapsto 0,\ h_j \mapsto 0), \\ \delta_2 &\mapsto R(\delta_2)\,(: t_1 \mapsto -1,\ t_2 \mapsto 1,\ h_j \mapsto 0), \\ \epsilon_i &\mapsto R(\epsilon_i)\,(: t_1, t_2 \mapsto 0,\ h_j \mapsto \delta_{ij}), \end{aligned}$$

is precisely the set of roots of $\mathfrak{g}(S_2, S_{4,2})$ relative to $\mathfrak{h}$.

Consider the lexicographic order on F with $\delta_1 > \delta_2 > \epsilon_1 > \epsilon_2 > \epsilon_3 > 0$. Then,

$$\Phi = \{\alpha_1 = \delta_1 - \epsilon_1 - \epsilon_2, \alpha_2 = 2\epsilon_2, \alpha_3 = \epsilon_1 - \epsilon_2 - \epsilon_3, \alpha_4 = 2\epsilon_3, \alpha_5 = \delta_2 - \epsilon_1\}$$

is a linearly independent set with $\Phi \subseteq \mathbb{Z}\Pi$, so that any positive element in Φ is a sum of elements in Π. Consider the elements:

$$\begin{array}{lll} E_1 = e_0^+ \otimes v_1 \otimes v_2, & H_1 = (t_1 - t_2) + h_1 + h_2, & F_1 = \xi_1 e_0^- \otimes w_1 \otimes w_2, \\ E_2 = \gamma_{w_2,w_2}, & H_2 = h_2, & F_2 = \xi_2 \gamma_{v_2,v_2}, \\ E_3 = w_1 \otimes v_2 \otimes v_3, & H_3 = -h_1 + h_2 + h_3, & F_3 = \xi_3 v_1 \otimes w_2 \otimes w_3, \\ E_4 = \gamma_{w_3,w_3}, & H_4 = h_3, & F_4 = \xi_4 \gamma_{v_3,v_3}, \\ E_5 = e_1^+ \otimes v_1, & H_5 = (t_1 - t_2) + h_1, & F_5 = \xi_5 e_1^- \otimes w_1, \end{array}$$

where $\xi_1, \dots, \xi_5$ are suitable scalars so as to have $[E_i, F_j] = \delta_{ij} H_j$ for any i, j.

With these elements $\mathfrak{g}(S_2, S_{4,2})$ is $\mathbb{Z}$-graded, by assining degree 1 to $E_1, \dots, E_5$, and degree -1 to $F_1, \dots, F_5$, and the hypotheses of Theorem 4.5 are satisfied relative to the rank 4 matrix (and Dynkin diagram):

$$A_{S_2,S_{4,2}} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 \\ -1 & 2 & -1 & 0 & 0 \\ 0 & -1 & 0 & -1 & 1 \\ 0 & 0 & -1 & 2 & 0 \\ 0 & 0 & 1 & 0 & 0 \end{pmatrix},$$



Therefore:

Proposition 5.15. *The Lie superalgebra* $\mathfrak{g}(S_2, S_{4,2})$ *is isomorphic to the centerless contragredient Lie superalgebra* $\mathfrak{g}\big(A_{S_2,S_{4,2}}, \{3, 5\}\big)/\mathfrak{c}$. *It is not simple, but its derived superalgebra* $[\mathfrak{g}(S_2, S_{4,2}), \mathfrak{g}(S_2, S_{4,2})]$ *is a simple ideal of codimension* 1*.*

Also,

$$\Pi_{\bar 0} = \{\beta_1 = 2\epsilon_1, \beta_2 = \delta_1 - \epsilon_1 - \epsilon_2, \beta_3 = 2\epsilon_2, \beta_4 = \delta_2 - \epsilon_2 - \epsilon_3, \beta_5 = 2\epsilon_3\}$$

is a linearly independent set with $\Phi_{\bar 0} \subseteq \mathbb{Z}\Pi_{\bar 0}$, so that any positive element in Φ is a sum of elements in $\Pi_{\bar 0}$, and with similar arguments one obtains that the associated Cartan matrix is of type A_5, $\mathfrak{g}_{\bar 0}$ thus being isomorphic to the centerless contragredient Lie algebra $\mathfrak{g}(A_5)/\mathfrak{c}$, which is isomorphic to the projective general Lie algebra $\mathfrak{pgl}_6$. The highest degree in $\mathfrak{g}_{\bar 1}$ is $\delta_1 + \delta_2 + \epsilon_3$, so the highest weight is $R(\delta_1 + \delta_2 + \epsilon_3) = \omega_3$. Therefore:

Proposition 5.16. $\mathfrak{g}(S_2, S_{4,2})$ *is a Lie superalgebra with even part isomorphic to* $\mathfrak{pgl}_6$*, and odd part isomorphic to the third exterior power of the natural module for* $\mathfrak{gl}_6$ *(of dimension* 20*).*

Since the characteristic is 3, this third exterior power is indeed a module for $\mathfrak{pgl}_6 = \mathfrak{gl}_6/k1$.

5.7. $\mathfrak{g}(S_{1,2}, S_{1,2})$. Now, let us consider two copies of $S_{1,2}$: $k1 \oplus V_i$, $i = 1, 2$ (Example 2.7). Then the even and odd parts of

$$\mathfrak{g} = \mathfrak{g}(S_{1,2}, S_{1,2}) = \bigl(\mathfrak{tri}(S_{1,2}) \oplus \mathfrak{tri}(S_{1,2})\bigr) \oplus \bigl(\oplus_{i=0}^2 \iota_i(S_{1,2} \otimes S_{1,2})\bigr)$$

are given, because of Corollary 2.12, by:

$$\begin{aligned} \mathfrak{g}_{\bar 0} &= \bigl(\mathfrak{sp}(V_1) \oplus \mathfrak{sp}(V_2)\bigr) \oplus \bigl(\oplus_{i=0}^2 k\iota_i(1 \otimes 1)\bigr) \oplus \bigl(\oplus_{i=0}^2 \iota_i(V_1 \otimes V_2)\bigr), \\ \mathfrak{g}_{\bar 1} &= \Bigl(V_1 \oplus \bigl(\oplus_{i=0}^2 \iota_i(V_1 \otimes 1)\bigr)\Bigr) \oplus \Bigl(V_2 \oplus \bigl(\oplus_{i=0}^2 \iota_i(1 \otimes V_2)\bigr)\Bigr). \end{aligned}$$

Consider the quaternion algebra

$$Q = k1 \oplus \bigl(\oplus_{i=0}^2 kx_i\bigr),$$

with $x_i^2 = -1$ for any $i = 0, 1, 2$, and $x_i x_{i+1} = -x_{i+1}x_i = x_{i+2}$ for any i (indices modulo 3). Its norm N satisfies $N(1) = N(x_i) = 1$, $i = 0, 1, 2$, and $N(1, x_i) = N(x_i, x_j) = 0$ for any $i \neq j$ ($N(a, b) = N(a + b) - N(a) - N(b)$).

Since the characteristic of k is 3, $N(1+x_1+x_2) = 0$, hence N represents 0 and Q is the split quaternion algebra. That is, Q is isomorphic to $\mathrm{Mat}_2(k)$. In particular, $Q_0 = (k1)^\perp = kx_0 \oplus kx_1 \oplus kx_2$ is a Lie algebra under the commutator $[x, y] = xy - yx$, which is isomorphic to $\mathfrak{sl}_2$.

The subspace $\oplus_{i=0}^2 k\iota_i(1 \otimes 1)$ in $\mathfrak{g}_{\bar 0}$ is a subalgebra, and

$$[\iota_i(1 \otimes 1), \iota_{i+1}(1 \otimes 1)] = \iota_{i+2}\bigl((1 \bullet 1) \otimes (1 \bullet 1)\bigr) = \iota_{i+2}(1 \otimes 1),$$

while $[x_i, x_{i+1}] = 2x_{i+2} = -x_{i+2}$. Therefore, $\oplus_{i=0}^2 k\iota_i(1 \otimes 1)$ is isomorphic to Q_0 under the linear map that takes $\iota_i(1 \otimes 1)$ to $-x_i$.

As vector spaces, there is a natural isomorphism

$$\Gamma_{\bar 0} : \mathfrak{g}_{\bar 0} \to \bigl(\mathfrak{sp}(V_1) \oplus \mathfrak{sp}(V_2) \oplus Q_0\bigr) \oplus \bigl(V_1 \otimes V_2 \otimes Q_0\bigr),$$

which is the identity on $\mathfrak{sp}(V_1) \oplus \mathfrak{sp}(V_2)$, and such that

$$\Gamma_{\bar 0}\bigl(\iota_i(1 \otimes 1)\bigr) = -x_i, \quad \Gamma_{\bar 0}\bigl(\iota_i(u_1 \otimes u_2)\bigr) = u_1 \otimes u_2 \otimes x_i,$$

for any $i = 0, 1, 2$ and $u_1 \in V_1$, $u_2 \in V_2$.

The Lie bracket in $\mathfrak{g}_{\bar 0}$ is then transfered to the right hand side as follows:

- $\mathfrak{sp}(V_1) \oplus \mathfrak{sp}(V_2) \oplus Q_0$ is a Lie subalgebra with componentwise bracket.
- $[s_1, u_1 \otimes u_2 \otimes p] = s_1(u_1) \otimes u_2 \otimes p$, $[s_2, u_1 \otimes u_2 \otimes p] = u_1 \otimes s_2(u_2) \otimes p$, for any $s_i \in \mathfrak{sp}(V_i)$, $u_i \in V_i$, $i = 1, 2$, and $p \in Q_0$.
- $[q, u_1 \otimes u_2 \otimes p] = u_1 \otimes u_2 \otimes [q, p]$, for any $u_i \in V_i$, $i = 1, 2$, and $p, q \in Q_0$. This is because $[\iota_i(1 \otimes 1), \iota_i(u_1 \otimes u_2)] = 0$, and $[\iota_i(1 \otimes 1), \iota_{i+1}(u_1 \otimes u_2)] = \iota_{i+2}(u_1 \otimes u_2)$ and $[\iota_i(1 \otimes 1), \iota_{i+2}(u_1 \otimes u_2)] = -\iota_{i+1}(u_1 \otimes u_2)$ in $\mathfrak{g}_{\bar 0}$ (see Section 3).
- For any $u_i, u_i' \in V_i$, $i = 1, 2$, and $p, p' \in Q_0$:

$$\begin{aligned} &[u_1 \otimes u_2 \otimes p, u_1' \otimes u_2' \otimes p'] \\ &\quad = -\langle u_2 | u_2'\rangle N(p, p')\gamma_{u_1, u_1'} - \langle u_1 | u_1'\rangle N(p, p')\gamma_{u_2, u_2'} - \langle u_1 | u_1'\rangle\langle u_2 | u_2'\rangle [p, q]. \end{aligned}$$

  This is because $[\iota_i(u_1 \otimes u_2), \iota_i(u_1' \otimes u_2')] = \langle u_2 | u_2'\rangle \gamma_{u_1, u_1'} + \langle u_1 | u_1'\rangle \gamma_{u_2, u_2'}$ (note that $S_{1,2} = k1 \oplus V$, and $\sigma_{u,u'} = -\gamma_{u,u'}$ (2.15)), $N(1, 1) = 2 = -1$ and $[\iota_i(u_1 \otimes u_2), \iota_{i+1}(u_1' \otimes u_2')] = -\iota_{i+2}\bigl((u_1 \bullet u_1') \otimes (u_2 \bullet u_2')\bigr) =$

$-\langle u_1|u_1'\rangle\langle u_2|u_2'\rangle\iota_{i+2}(1\otimes 1)$, which corresponds to $[u_1\otimes u_2\otimes x_i, u_1'\otimes u_2'\otimes x_{i+1}] = \langle u_1|u_1'\rangle\langle u_2|u_2'\rangle x_{i+2}$ ($[x_i,x_{i+1}] = -x_{i+2}$).

Now, as vector spaces, there is also a natural isomorphism:

$$\begin{aligned}\Gamma_{\bar 1}: \mathfrak{g}_{\bar 1} &\longrightarrow (V_1\oplus V_2)\otimes Q,\\ u_1 &\mapsto\ u_1\otimes 1,\\ u_2 &\mapsto\ -u_2\otimes 1,\\ \iota_i(u_1\otimes 1) &\mapsto -u_1\otimes x_i,\\ \iota_i(1\otimes u_2) &\mapsto u_2\otimes x_i,\end{aligned}$$

for $i=0,1,2$, $u_1\in V_1$, and $u_2\in V_2$. Under $\Gamma_{\bar 0}$ and $\Gamma_{\bar 1}$, the Lie bracket $\mathfrak{g}_{\bar 0}\times\mathfrak{g}_{\bar 1}\to\mathfrak{g}_{\bar 1}$ is transferred to:

- $[s_i,(u_1+u_2)\otimes q] = s_i(u_i)\otimes q$ for any $s_i\in\mathfrak{sp}(V_i)$, $u_i\in V_i$, $i=1,2$ and $q\in Q$.
- $[p,(u_1+u_2)\otimes q] = (u_1+u_2)\otimes pq$, for any $p\in Q_0$, $q\in Q$, and $u_i\in V_i$, $i=1,2$.
- $[u_1\otimes u_2\otimes p, u_1'\otimes q] = -\langle u_1|u_1'\rangle u_2\otimes pq$, for any $u_1,u_1'\in V_1$, $u_2\in V_2$, $p\in Q_0$ and $q\in Q$.
- $[u_1\otimes u_2\otimes p, u_2'\otimes q] = \langle u_2|u_2'\rangle u_1\otimes pq$, for any $u_1\in V_1$, $u_2,u_2'\in V_2$, $p\in Q_0$ and $q\in Q$.

While the bracket $\mathfrak{g}_{\bar 1}\times\mathfrak{g}_{\bar 1}\to\mathfrak{g}_{\bar 0}$ transfers to:

- $[u_1\otimes q, u_1'\otimes q'] = -N(q,q')\gamma_{u_1,u_1'} - \langle u_1,u_1'\rangle(q\bar q' - q'\bar q)$,
- $[u_2\otimes q, u_2'\otimes q'] = -N(q,q')\gamma_{u_2,u_2'} - \langle u_2,u_2'\rangle(q\bar q' - q'\bar q)$,
- $[u_1\otimes q, u_2\otimes q'] = -u_1\otimes u_2\otimes(q\bar q' - q'\bar q)$,

for any $u_i,u_i'\in V_i$, $i=1,2$, and $q,q'\in Q$. Here $^-$ denotes the canonical involution in the quaternion algebra Q (that is, the symplectic involution in $\mathrm{Mat}_2(k)$).

Therefore, the even and odd parts of $\mathfrak{g}=\mathfrak{g}(S_{1,2},S_{1,2})$ can be identified to

$$\begin{aligned}\mathfrak{g}_{\bar 0} &= \bigl(\mathfrak{sp}(V_1)\oplus\mathfrak{sp}(V_2)\oplus\mathfrak{sl}_2\bigr)\oplus\bigl(V_1\otimes V_2\otimes\mathfrak{sl}_2\bigr),\\ \mathfrak{g}_{\bar 1} &= (V_1\oplus V_2)\otimes\mathfrak{gl}_2\,.\end{aligned}\tag{5.17}$$

The action of $\mathfrak{sl}_2$ on $\mathfrak{gl}_2$ is given by left multiplication, hence $\mathfrak{gl}_2$ decomposes, as a module for $\mathfrak{sl}_2$, as $\mathfrak{gl}_2 = \mathfrak{j}_1\oplus\mathfrak{j}_{-1}$, where $\mathfrak{j}_1$ (respectively $\mathfrak{j}_{-1}$) consists of the 2×2 matrices with zero second (resp. first) column.

Let $\{v_i,w_i\}$ be a symplectic basis of V_i, $i=1,2$, as considered so far, and let $\{h=E_{11}-E_{22}, e=E_{12}, f=E_{21}\}$ be the standard basis of $\mathfrak{sl}_2$ (E_{ij} denotes the 2×2 matrix with entry (i,j) equal to 1 and all the other entries 0). Note that $\{E_{11},E_{21}\}$ is a basis of $\mathfrak{j}_1$, while $\{E_{12},E_{22}\}$ is a basis of $\mathfrak{j}_{-1}$.

Consider here the free $\mathbb{Z}$-module F with basis $\epsilon_1,\epsilon_2,\epsilon,\delta$. The Lie superalgebra $\mathfrak{g}=\mathfrak{g}(S_{1,2},S_{1,2})$ is F-graded by assigning degree ϵ_i to w_i and $-\epsilon_i$ to v_i as usual, $i=1,2$, degree $\epsilon+\delta$ to E_{11} ($\in\mathfrak{j}_1$), $-\epsilon+\delta$ to E_{21} ($\in\mathfrak{j}_1$), $\epsilon-\delta$ to E_{12} ($\in\mathfrak{j}_{-1}$), and $-\epsilon-\delta$ to E_{22} ($\in\mathfrak{j}_{-1}$). Then it follows that, for instance, the degree of $v_1\otimes E_{22}$ is $-\epsilon_1-\epsilon-\delta$, or that the degree of $e=E_{12}\in\mathfrak{sl}_2$ is 2ϵ.

The sets of nonzero even and odd degrees are:

$$\begin{aligned}
\Phi_{\bar 0} &= \{\pm 2\epsilon_1, \pm 2\epsilon_2, \pm 2\epsilon, \pm\epsilon_1 \pm \epsilon_2, \pm\epsilon_1 \pm \epsilon_2 \pm 2\epsilon\},\\
\Phi_{\bar 1} &= \{\pm\epsilon_1 \pm \epsilon \pm \delta, \pm\epsilon_2 \pm \epsilon \pm \delta\}.
\end{aligned}$$

The abelian subalgebra $\mathfrak{h} = kh_1 \oplus kh_2 \oplus kh$ ($h_i = \gamma_{v_i,w_i} \in \mathfrak{sp}(V_i)$, $i = 1, 2$, as usual) is a Cartan subalgebra of $\mathfrak{g}$ and the image of $\Phi_{\bar 0} \cup \Phi_{\bar 1}$ under the homomorphism of abelian groups:

$$\begin{aligned}
R : F &\longrightarrow \mathfrak{h}^*\\
\epsilon_1 &\mapsto R(\epsilon_1)\,(: h_1 \mapsto 1,\ h_2 \mapsto 0,\ h \mapsto 0),\\
\epsilon_2 &\mapsto R(\epsilon_2)\,(: h_1 \mapsto 0,\ h_2 \mapsto 1,\ h \mapsto 0),\\
\epsilon &\mapsto R(\epsilon)\,(: h_1 \mapsto 0,\ h_2 \mapsto 0,\ h \mapsto 1),\\
\delta &\mapsto 0,
\end{aligned}$$

is precisely the set of roots of $\mathfrak{g}$ relative to $\mathfrak{h}$.

Consider the lexicographic order on F with $\delta > \epsilon_1 > \epsilon_2 > \epsilon > 0$, then

$$\Phi = \{\alpha_1 = 2\epsilon_2, \alpha_2 = \epsilon_1 - \epsilon_2 - 2\epsilon, \alpha_3 = 2\epsilon, \alpha_4 = \delta - \epsilon_1 - \epsilon\}$$

is a linearly independent set in F with $\Phi = \Phi_{\bar 0} \cup \Phi_{\bar 1} \subseteq \mathbb{Z}\Pi$, so that any positive element in Φ is a sum of elements in Π. Consider the elements:

$$\begin{aligned}
&E_1 = \gamma_{w_2,w_2}, &&E_2 = w_1 \otimes v_2 \otimes E_{21}, &&E_3 = e, &&E_4 = v_1 \otimes E_{21},\\
&F_1 = -\gamma_{v_2,v_2}, &&F_2 = -v_1 \otimes w_2 \otimes E_{12}, &&F_3 = f, &&F_4 = w_1 \otimes E_{12}.
\end{aligned}$$

Then:

$$\begin{aligned}
H_1 &= [E_1, F_1] = \gamma_{v_2,w_2} = h_2,\\
H_2 &= [E_2, F_2] = -\gamma_{v_1,w_1} + \gamma_{v_2,w_2} + h = -h_1 + h_2 + h,\\
H_3 &= [E_3, F_3] = h,\\
H_4 &= [E_4, F_4] = \gamma_{v_1,w_1} - h = h_1 - h,
\end{aligned}$$

and the hypotheses of Theorem 4.7 are satisfied relative to the rank 3 matrix (and Dynkin diagram):

$$A_{S_{1,2},S_{1,2}} = \begin{pmatrix} 2 & -1 & 0 & 0\\ -1 & 2 & -1 & 0\\ 0 & -2 & 2 & -1\\ 0 & 0 & 1 & 0 \end{pmatrix},$$



Therefore:

Proposition 5.18. *The Lie superalgebra* $\mathfrak{g}(S_{1,2}, S_{1,2})$ *is isomorphic to the centerless derived contragredient Lie superalgebra* $\mathfrak{g}'\big(A_{S_{1,2},S_{1,2}}, \{4\}\big)/\mathfrak{c}$.

Moreover, $\{E_i, F_i, H_i : i = 1, 2, 3\}$ generate the even part of $\mathfrak{g}_{\bar 0}$, the associated Cartan subalgebra is of type B_3, $\mathfrak{g}_{\bar 0}$ thus being isomorphic to the orthogonal Lie algebra $\mathfrak{so}_7$. The grading of $\mathfrak{g}$ by F gives, in particular, a consistent $\mathbb{Z}$-grading by means of the projection $F = \mathbb{Z}\epsilon_1 \oplus \mathbb{Z}\epsilon_2 \oplus \mathbb{Z}\epsilon \oplus \mathbb{Z}\delta \to \mathbb{Z}$, $n_1\epsilon_1 + n_2\epsilon_2 + n_3\epsilon + n_4\delta \mapsto n_4$, where

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_1,$$

with

$$\mathfrak{g}_0 = \mathfrak{g}_{\bar 0}, \quad \mathfrak{g}_1 = (V_1 \oplus V_2) \otimes \mathfrak{j}_1, \quad \mathfrak{g}_{-1} = (V_1 \oplus V_2) \otimes \mathfrak{j}_{-1}\,.$$

As modules over $\mathfrak{g}_{\bar 0} = \mathfrak{g}_0$, $\mathfrak{g}_1$ and $\mathfrak{g}_{-1}$ are isomorphic, and $\mathfrak{g}_1$ is an irreducible module with highest weight $R(\epsilon_1 + \epsilon + \delta) = \omega_3$, so $\mathfrak{g}_1$ is the spin module for $\mathfrak{g}_0$. Hence:

Proposition 5.19. *$\mathfrak{g}(S_{1,2}, S_{1,2})$ is a simple Lie superalgebra whose even part is isomorphic to $\mathfrak{so}_7$, and its odd part is the direct sum of two copies of the spin module for the even part.*

Corollary 5.20. *The Lie superalgebra $\mathfrak{g}(S_{1,2}, S_{1,2})$ is isomorphic to the Lie superalgebra in* [Eld06b, Theorem 4.23(ii)], *constructed in terms of a null orthogonal triple system.*

5.8. $\mathfrak{g}(S_1, S_{1,2})$. The Lie superalgebra $\mathfrak{g}(S_1, S_{1,2})$ is a subsuperalgebra of $\mathfrak{g}(S_{1,2}, S_{1,2})$. Given the description of this latter superalgebra in 5.7, more specifically in (5.17), the even and odd parts of $\mathfrak{g} = \mathfrak{g}(S_1, S_{1,2})$ can be described as

$$\begin{aligned} \mathfrak{g}_{\bar 0} &= \mathfrak{sp}(V_2) \oplus \mathfrak{sl}_2 \ (\subseteq \mathfrak{g}(S_{1,2}, S_{1,2})_{\bar 0}), \\ \mathfrak{g}_{\bar 1} &= V_2 \otimes \mathfrak{gl}_2 \ (\subseteq \mathfrak{g}(S_{1,2}, S_{1,2})_{\bar 1}), \end{aligned}$$

with

$$\Phi_{\bar 0} = \{\pm 2\epsilon_2, \pm 2\epsilon\}, \quad \Phi_{\bar 1} = \{\pm\epsilon_2 \pm \epsilon \pm \delta\}.$$

With the lexicographic order in which $\delta > \epsilon_2 > \epsilon > 0$,

$$\Pi = \{\alpha_1 = 2\epsilon_2, \alpha_2 = \delta - \epsilon_2 - \epsilon, \alpha_3 = 2\epsilon\},$$

with associated matrix and Dynkin diagram:

$$A_{S_1,S_{1,2}} = \begin{pmatrix} 2 & -1 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 2 \end{pmatrix},$$



and the same arguments as in the previous section give:

Proposition 5.21. *The Lie superalgebra $\mathfrak{g}(S_1, S_{1,2})$ is isomorphic to the centerless derived contragredient Lie superalgebra $\mathfrak{g}'\big(A_{S_1,S_{1,2}}, \{2\}\big)/\mathfrak{c}$.*

Corollary 5.22. *$\mathfrak{g}(S_1, S_{1,2})$ is isomorphic to the projective special Lie superalgebra $\mathfrak{psl}(2,2)$.*

This is the only Lie superalgebra in the Extended Freudenthal's Magic Square with a counterpart in characteristic 0.

Note that the even part $\mathfrak{g}_{\bar 0}$ is the direct sum of two copies of $\mathfrak{sl}_2$, while the odd part is a direct sum of two copies of the four dimensional irreducible module for $\mathfrak{g}_{\bar 0}$ which consists of the tensor product of the natural modules for each copy of $\mathfrak{sl}_2$.

5.9. **$\mathfrak{g}(S_4, S_{1,2})$.** Here a description in terms of $V(\sigma)$'s is again possible. Three copies of V are needed for S_4 and an extra copy for $S_{1,2}$. The indices 1, 2 and 3 will be used for the three copies related to S_4 and the index 4 for the extra copy attached to $S_{1,2}$. Then

$$\begin{aligned}\mathfrak{g}(S_4, S_{1,2}) &= \bigl(\mathfrak{tri}(S_4) \oplus \mathfrak{tri}(S_{1,2})\bigr) \oplus \bigl(\oplus_{i=0}^{2} \iota_i(S_4 \otimes S_{1,2})\bigr)\\ &= \oplus_{\sigma \in \mathcal{S}_{S_4, S_{1,2}}} V(\sigma),\end{aligned}$$

where

$$\mathcal{S}_{S_4,S_{1,2}} = \bigl\{\emptyset, \{4\}, \{1,2\}, \{1,2,4\}, \{2,3\}, \{2,3,4\}, \{1,3\}, \{1,3,4\}\bigr\}.$$

The nonzero even and odd degrees are:

$$\begin{aligned}\Phi_{\bar 0} &= \{\pm 2\epsilon_i : 1 \le i \le 4\} \cup \{\pm\epsilon_1 \pm \epsilon_2, \pm\epsilon_2 \pm \epsilon_3, \pm\epsilon_1 \pm \epsilon_3\},\\ \Phi_{\bar 1} &= \{\pm\epsilon_4, \pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_4, \pm\epsilon_2 \pm \epsilon_3 \pm \epsilon_4, \pm\epsilon_1 \pm \epsilon_3 \pm \epsilon_4\}.\end{aligned}$$

With the lexicographic order given by $0 < \epsilon_1 < \epsilon_2 < \epsilon_3 < \epsilon_4$, the set of irreducible degrees is

$$\Pi = \{\alpha_1 = \epsilon_3 - \epsilon_2, \alpha_2 = \epsilon_2 - \epsilon_1, \alpha_3 = 2\epsilon_1, \alpha_4 = \epsilon_4 - \epsilon_2 - \epsilon_3\},$$

which is a linearly independent set over $\mathbb{Z}$, and $\Phi = \Phi_{\bar 0} \cup \Phi_{\bar 1}$ is contained in $\mathbb{Z}\Pi$. The associated matrix and Dynkin diagram are:

$$A_{S_4,S_{1,2}} = \begin{pmatrix} 2 & -1 & 0 & 0\\ -1 & 2 & -2 & -1\\ 0 & -1 & 2 & 0\\ 0 & 1 & 0 & 0 \end{pmatrix},$$



and

Proposition 5.23. *The Lie superalgebra $\mathfrak{g}(S_4, S_{1,2})$ is isomorphic to the contragredient Lie superalgebra $\mathfrak{g}\bigl(A_{S_4,S_{1,2}}, \{4\}\bigr)$.*

Also, the set of irreducible even degrees is

$$\Pi_{\bar 0} = \{\alpha_1, \alpha_2, \alpha_3, \tilde\alpha_4 = 2\epsilon_4\},$$

with associated Cartan matrix of type $C_3 \oplus A_1$. (This also follows from $\mathfrak{g}(S_4, S_{1,2})_{\bar 0} = \mathfrak{g}(S_4, S_1) \oplus \mathfrak{sp}(V_4)$.)

The odd part, which has dimension 26, is irreducible with highest weight $R(\epsilon_2 + \epsilon_3 + \epsilon_4) = R(\epsilon_2 + \epsilon_3) + R(\epsilon_4)$, so that:

Proposition 5.24. *$\mathfrak{g}(S_4, S_{1,2})$ is a simple Lie superalgebra with even part isomorphic to the direct sum $\mathfrak{sp}_6 \oplus \mathfrak{sl}_2$, and odd part the irreducible module which is the tensor product of the irreducible module $V(\omega_2)$ for $\mathfrak{sp}_6$ (of dimension 13) , and the natural two dimensional module for $\mathfrak{sl}_2$.*

The irreducible module $V(\omega_2)$ for $\mathfrak{sp}_6$ can be described as follows. Let W be the natural six-dimensional module for $\mathfrak{sp}_6$, which is endowed with a nondegenerate alternating form $\{.|.\}$, and take a symplectic basis $\{a_i, b_i : i = 1,2,3\}$. Consider the linear map $\varphi : \bigwedge^2 W, \to k,\ z_1 \wedge z_2 \mapsto \{z_1|z_2\}$. Then, since the characteristic of k is 3, $k(\sum_{i=1}^{3} a_i \wedge b_i)$ is a trivial submodule of $\ker\varphi$, and $\ker\varphi / k(\sum_{i=1}^{3} a_i \wedge b_i)$ is an irreducible module with highest weight vector the class of $b_2 \wedge b_3$ modulo $k(\sum_{i=1}^{3} a_i \wedge b_i)$ (conventions as in Subsection 5.1). This is, up to isomorphism, the required irreducible module.

5.10. $\mathfrak{g}(S_8, S_{1,2})$**.** Here the indices 1, 2, 3 and 4 will be used for the copies of V attached to S_8, while the index 5 will be reserved for the copy of V attached to $S_{1,2}$. With the same arguments as before:

$$\begin{aligned}\mathfrak{g}(S_8, S_{1,2}) &= \bigl(\mathfrak{tri}(S_8) \oplus \mathfrak{tri}(S_{1,2})\bigr) \oplus \bigl(\oplus_{i=0}^{2} \iota_i(S_8 \otimes S_{1,2})\bigr)\\ &= \oplus_{\sigma \in \mathcal{S}_{S_8,S_{1,2}}} V(\sigma),\end{aligned}$$

where

$$\begin{aligned}\mathcal{S}_{S_8,S_{1,2}} = \bigl\{&\emptyset, \{1,2,3,4\}, \{5\},\\ &\{1,2\}, \{3,4\}, \{1,2,5\}, \{3,4,5\},\\ &\{2,3\}, \{1,4\}, \{2,3,5\}, \{1,4,5\},\\ &\{1,3\}, \{2,4\}, \{1,3,5\}, \{2,4,5\}\bigr\}.\end{aligned}$$

Here,

$\Phi_{\bar{0}} = \{\pm 2\epsilon_i : 1 \leq i \leq 5\} \cup \{\pm\epsilon_1 \pm \epsilon_2 \pm \epsilon_3 \pm \epsilon_4\} \cup \{\pm\epsilon_i \pm \epsilon_j : 1 \leq i < j \leq 4\}$,
$\Phi_{\bar{1}} = \{\pm\epsilon_5\} \cup \{\pm\epsilon_i \pm \epsilon_j \pm \epsilon_5 : 1 \leq i < j \leq 4\}$,

and with the lexicographic order given by $0 < \epsilon_1 < \epsilon_2 < \epsilon_3 < \epsilon_4 < \epsilon_5$, the set of irreducible degrees is

$$\begin{aligned}\Pi = \{\alpha_1 = \epsilon_4 - \epsilon_3 - \epsilon_2 - \epsilon_1, \alpha_2 = \epsilon_1, \alpha_3 = \epsilon_2 - \epsilon_1,&\\ \alpha_4 = \epsilon_3 - \epsilon_2, \alpha_5 = \epsilon_5 - \epsilon_3 - \epsilon_4\}&,\end{aligned}$$

which is a $\mathbb{Z}$-linearly independent set with $\Phi \subseteq \mathbb{Z}\Pi$. The associated matrix and Dynkin diagrams are

$$A_{S_8,S_{1,2}} = \begin{pmatrix} 2 & -1 & 0 & 0 & 0\\ -1 & 2 & -1 & 0 & 0\\ 0 & -2 & 2 & -1 & 0\\ 0 & 0 & -1 & 2 & -1\\ 0 & 0 & 0 & 1 & 0 \end{pmatrix},$$



and then:

Proposition 5.25. *The Lie superalgebra* $\mathfrak{g}(S_8, S_{1,2})$ *is isomorphic to the contragredient Lie superalgebra* $\mathfrak{g}\bigl(A_{S_8,S_{1,2}}, \{5\}\bigr)$.

Also,

$$\Pi_{\bar{0}} = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \tilde{\alpha}_5 = 2\epsilon_5\},$$

with associated Cartan matrix of type $F_4 \oplus A_1$ (this also follows from $\mathfrak{g}(S_8, S_{1,2})_{\bar{0}} = \mathfrak{g}(S_8, S_1) \oplus \mathfrak{sp}(V_5)$). The odd part, which has dimension 50, is irreducible with highest weight $R(\epsilon_5 + \epsilon_4 + \epsilon_3) = R(\epsilon_3 + \epsilon_4) + R(\epsilon_5)$. $R(\epsilon_5)$ is the highest weight of the natural two dimensional module for $\mathfrak{sp}(V_5) \cong \mathfrak{sl}_2$, while $R(\epsilon_3 + \epsilon_4)$ is the highest weight ω_4 for the direct summand of $\mathfrak{g}(S_8, S_{1,2})_{\bar{0}}$ isomorphic to $\mathfrak{f}_4$, the exceptional simple split Lie algebra of type F_4 (with the ordering of the roots given above). Thus:

Proposition 5.26. *$\mathfrak{g}(S_8, S_{1,2})$ is a simple Lie superalgebra with even part isomorphic to $\mathfrak{f}_4 \oplus \mathfrak{sl}_2$, and odd part the irreducible module which is the tensor product of the natural two dimensional module for $\mathfrak{sl}_2$ and the 25-dimensional module of highest weight ω_4 for $\mathfrak{f}_4$.*

In characteristic > 3, the module $V(\omega_4)$ for $\mathfrak{f}_4$ is usually described as the set of trace zero elements J_0 in the split exceptional Jordan algebra $J = H_3(C)$, where C is the split octonion algebra. However, in characteristic 3, the identity matrix belongs to J_0, and $J_0/k1$ is the irreducible module $V(\omega_4)$ in this case.

5.11. **$\mathfrak{g}(S_2, S_{1,2})$.** Take the basis $\{e^+, e^-\}$ of the symmetric composition algebra S_2 as in 5.6, and identify $\mathfrak{tri}(S_2)$ with $\mathfrak{t} = \text{span}\{t_1, t_2\}$ as was done there. Take also three copies S_2^i (with basis $\{e_i^+, e_i^-\}$) of S_2, $i = 0, 1, 2$. Here we consider the Lie superalgebra

$$\mathfrak{g} = \mathfrak{g}(S_2, S_{1,2}) = \bigl(\mathfrak{tri}(S_2) \oplus \mathfrak{tri}(S_{1,2})\bigr) \oplus \bigl(\oplus_{i=0}^{2} \iota_i(S_2 \otimes S_{1,2})\bigr),$$

so, with standard identifications:

$$\begin{aligned} \mathfrak{g}_{\bar 0} &= \bigl(\mathfrak{t} \oplus \mathfrak{sp}(V)\bigr) \oplus \bigl(\oplus_{i=0}^{2} S_2^i\bigr), \\ \mathfrak{g}_{\bar 1} &= V \oplus \bigl(\oplus_{i=0}^{2} (S_2^i \otimes V)\bigr). \end{aligned}$$

Consider the free $\mathbb{Z}$-module F freely generated by δ_1, δ_2 and ϵ. With the conventions in 5.6, $\mathfrak{g}$ is F-graded and the sets of nonzero even and odd degrees are:

$$\begin{aligned} \Phi_{\bar 0} &= \{\pm 2\epsilon, \pm\delta_1, \pm\delta_2, \pm(\delta_1 + \delta_2)\} \\ \Phi_{\bar 1} &= \{\pm\epsilon, \pm\delta_1 \pm \epsilon, \pm\delta_2 \pm \epsilon, \pm(\delta_1 + \delta_2) \pm \epsilon\}. \end{aligned}$$

With the lexicographic order in which $\epsilon > \delta_1 > \delta_2 > 0$,

$$\Pi = \{\alpha_1 = \delta_1, \alpha_2 = \delta_2, \alpha_3 = \epsilon - (\delta_1 + \delta_2)\}$$

is a linearly independent set with $\Phi = \Phi_{\bar 0} \cup \Phi_{\bar 1} \subseteq \mathbb{Z}\Pi$. Now consider the elements ($\{v, w\}$ denotes a symplectic basis of V):

$$\begin{aligned} &E_1 = e_0^+ (= \iota_0(1 \otimes e^+)), && E_2 = e_1^+, && E_3 = e_2^+ \otimes w (= \iota_2(e^+ \otimes w)), \\ &H_1 = 2(t_1 - t_2), && H_2 = 2(t_1 - t_2), && H_3 = (t_1 - t_2) - \gamma_{v,w}, \\ &F_1 = \xi_1 e_0^-, && F_2 = \xi_2 e_1^-, && F_3 = \xi_3 e_2^- \otimes v, \end{aligned}$$

where ξ_1, ξ_2, ξ_3 are suitable scalars so as to have $[E_i, F_j] = \delta_{ij} H_i$ for any i, j.

With these elements, $\mathfrak{g}(S_2, S_{1,2})$ is $\mathbb{Z}$-graded, by assigning degree 1 to E_1, E_2, E_3 and degree -1 to F_1, F_2, F_3, and the hypotheses of Theorem 4.5 are satisfied relative to the rank 2 matrix (with associated Dynkin diagram):

$$A_{S_2, S_{1,2}} = \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ 1 & 1 & 0 \end{pmatrix},$$



Therefore:

Proposition 5.27. *The Lie superalgebra $\mathfrak{g}(S_2, S_{1,2})$ is isomorphic to the centerless contragredient Lie superalgebra $\mathfrak{g}\bigl(A_{S_2,S_{1,2}}, \{3\}\bigr)/\mathfrak{c}$. It is not simple, but its derived superalgebra is a simple ideal of codimension 1.*

Also,

$$\Pi_{\bar{0}} = \{\beta_1 = \delta_1, \beta_2 = \delta_2, \beta_3 = 2\epsilon\}$$

is a linearly independent set with $\Phi_{\bar{0}} \subseteq \mathbb{Z}\Pi_{\bar{0}}$ and, with already used arguments, it is obtained that the associated Cartan matrix is of type $A_2 \oplus A_1$, $\mathfrak{g}_{\bar{0}}$ thus being isomorphic to the direct sum of the centerless contragredient Lie algebra $\mathfrak{g}(A_2)/\mathfrak{c}$, which is isomorphic to $\mathfrak{pgl}_3$, and of $\mathfrak{sl}_2$. The highest degree in $\mathfrak{g}_{\bar{1}}$ is $(\delta_1 + \delta_2) + \epsilon$, and $\mathfrak{g}_{\bar{1}}$ is easily seen to be irreducible. Hence:

Proposition 5.28. *$\mathfrak{g}(S_2, S_{1,2})$ is a Lie superalgebra with even part isomorphic to the direct sum of $\mathfrak{pgl}_3$ and $\mathfrak{sl}_2$, and odd part isomorphic, as a module for the even part, to the tensor product of the "adjoint module" $\mathfrak{psl}_3$ of $\mathfrak{pgl}_3$ by the natural two dimensional module for $\mathfrak{sl}_2$.*

Table 4 summarizes the information on the even and odd parts of the Lie superalgebras in the extended Freudenthal Magic Square obtained in this section. The notation (n) will indicate a module of dimension n.

	$S_{1,2}$	$S_{4,2}$
S_1	$\mathfrak{psl}_{2,2}$	$\mathfrak{sp}_6 \oplus (14)$
S_2	$\big(\mathfrak{pgl}_3 \oplus \mathfrak{sl}_2\big) \oplus \big(\mathfrak{psl}_3 \otimes (2)\big)$	$\mathfrak{pgl}_6 \oplus (20)$
S_4	$\big(\mathfrak{sp}_6 \oplus \mathfrak{sl}_2\big) \oplus \big((13) \otimes (2)\big)$	$\mathfrak{so}_{12} \oplus spin_{12}$
S_8	$\big(\mathfrak{f}_4 \oplus \mathfrak{sl}_2\big) \oplus \big((25) \otimes (2)\big)$	$\mathfrak{e}_7 \oplus (56)$
$S_{1,2}$	$\mathfrak{so}_7 \oplus 2spin_7$	$\mathfrak{sp}_8 \oplus (40)$
$S_{4,2}$		$\mathfrak{so}_{13} \oplus spin_{13}$

TABLE 4. Even and odd parts

With the exception of $\mathfrak{psl}_{2,2}$, none of these Lie superalgebras have a counterpart in characteristic 0. Corollary 5.11 shows that $\mathfrak{g}(S_{4,2}, S_{4,2})$ has already appeared in [Elda, Theorem 3.1(ii)], while Corollary 5.20 shows that $\mathfrak{g}(S_{1,2}, S_{1,2})$ is the Lie superalgebra that appears in [Eld06b, Theorem 4.23(ii)].

In a forthcoming paper, it will be shown that the Lie superalgebras $\mathfrak{g}(S_r, S_{1,2})$ and $\mathfrak{g}(S_r, S_{4,2})$, for $r = 1$, 4 and 8, and their derived subalgebras for $r = 2$, are precisely the simple Lie superalgebras defined in [Eld06b] in terms of symplectic and orthogonal triple systems related to simple Jordan algebras of degree 3.

REFERENCES

[AF93] B. N. Allison and J. R. Faulkner, *Nonassociative coefficient algebras for Steinberg unitary Lie algebras*, J. Algebra **161** (1993), no. 1, 1–19.

[Bae02] John C. Baez, *The octonions*, Bull. Amer. Math. Soc. (N.S.) **39** (2002), no. 2, 145–205.

[BS] C. H. Barton and A. Sudbery, *Magic Squares of Lie Algebras*, arXiv: math.RA/0001083.

[BS03] C. H. Barton and A. Sudbery, *Magic squares and matrix models of Lie algebras*, Adv. Math. **180** (2003), no. 2, 596–647.

[Car38] Elie Cartan, *Lecons sur la théorie des spineurs.*, Paris: Hermann & Cie. 96 p., 1938.

[Cun06] Isabel Cunha, Doctoral Thesis, in preparation.

[Eld04] Alberto Elduque, *The magic square and symmetric compositions.*, Rev. Mat. Iberoam. **20** (2004), no. 2, 475–491.

[Eld06a] ———, *A new look at Freudenthal's magic square*, Non-Associative Algebra and Its Applications, L. Sabinin, L. Sbitneva, and I.P. Shestakov, eds., Lecture Notes in Pure and Applied Mathematics, vol. 246, CRC Press, Taylor and Francis Group, Boca Raton, FL, 2006, pp. 149–165.

[Eld06b] ———, *New simple Lie superalgebras in characteristic 3*, J. Algebra **296** (2006), 196–233.

[Elda] ———, *Some new simple modular Lie superalgebras*, arXiv: math.RA/0512654. To appear in Pacific J. Math.

[Eldb] ———, *The Magic Square and Symmetric Compositions II*, arXiv: math.RT/0507282. To appear in Rev. Mat. Iberoamericana.

[EO02] Alberto Elduque and Susumu Okubo, *Composition superalgebras.*, Commun. Algebra **30** (2002), no. 11, 5447–5471.

[Fre64] H. Freudenthal, *Lie groups in the foundations of geometry*, Adv. Math. **1** (1964), 145–190.

[Hum72] J.E. Humphreys, *Introduction to Lie algebras and representation theory.*, Graduate Texts in Mathematics. 9. New York-Heidelberg-Berlin: Springer-Verlag. xii, 169 p., 1972.

[Kac77] V.G. Kac, *Lie superalgebras.*, Adv. Math. **26** (1977), 8–96.

[Kac90] ———, *Infinite dimensional Lie algebras. 3rd ed.*, Cambridge University Press. xxi, 400 p., 1990.

[KMRT98] Max-Albert Knus, Alexander Merkurjev, Markus Rost, and Jean-Pierre Tignol, *The book of involutions. With a preface by J. Tits.*, Colloquium Publications. American Mathematical Society (AMS). 44. Providence, RI: American Mathematical Society (AMS). xxi, 593 p., 1998.

[LM02] J. M. Landsberg and L. Manivel, *Triality, exceptional Lie algebras and Deligne dimension formulas*, Adv. Math. **171** (2002), no. 1, 59–85.

[LM04] ———, *Representation theory and projective geometry*, Algebraic transformation groups and algebraic varieties, Encyclopaedia Math. Sci., vol. 132, Springer, Berlin, 2004, pp. 71–122.

[Sch66] R.D. Schafer, *An introduction to nonassociative algebras*, Pure and Applied Mathematics, 22. A Series of Monographs and Textbooks. New York and London: Academic Press. X, 166 p., 1966.

[She97] I. P. Shestakov, *Prime alternative superalgebras of arbitrary characteristic*, Algebra i Logika **36** (1997), no. 6, 675–716, 722.

[Tit66] Jacques Tits, *Algèbres alternatives, algèbres de Jordan et algèbres de Lie exceptionnelles. I: Construction*, Nederl. Akad. Wet., Proc., Ser. A **69** (1966), 223–237.

[Vin05] E.B. Vinberg, *Construction of the exceptional simple Lie algebras.*, Vinberg, Ernest (ed.), Lie groups and invariant theory. Providence, RI: American Mathematical Society (AMS). Translations. Series 2. American Mathematical Society 213. Advances in the Mathematical Sciences 56, 241-242 , 2005.

[VK71] B.Yu. Vejsfejler and V.G. Kats, *Exponentials in Lie algebras of characteristic p.*, Izv. Akad. Nauk SSSR, Ser. Mat. **35** (1971), 762–788.

Departamento de Matemática, Universidade da Beira Interior,
6200 Covilhá, Portugal
E-mail address: `icunha@mat.ubi.pt`

Departamento de Matemáticas, Universidad de Zaragoza, 50009 Zaragoza, Spain
E-mail address: `elduque@unizar.es`